

Premises Address

(1). Syngenta India Limited, Santa Monica Works, Corlim, Ilhas,
Goa
Goa

(2). Unit II, Plot No. 155/1, GIDC Estate, Ankaleshwar, District-
Bharuch, Gujarat
Gujarat

(3). Plot No. 1, Swastik Ceramic Compound, AT & PO Sankarda,
District Baroda, Gujarat
Gujarat

Customer Care Details

Address: Syngenta India Limited, Amar Paradigm, S.No. 110/11/3,
Baner Road, Pune- 411 045,
Email: Rajesh.Dhawan@syngenta.com,
Contact No. 011-71010212

Safety Pictograms:



© 2011 Syngenta

(1).

(2).

(3).

»—« ± μ

© 2011 Syngenta India Limited, Amar Paradigm, S.No. 110/11/3, Baner
Road, Pune- 411 045,
p² μ -Rajesh.Dhawan@syngenta.com,
Contact No. 011-71010212

Secondary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Azoxystrobin 18.2% w/w Difenoconazole 11.4% w/w SC

BRAND NAME: -

Crops and Pest Groups: Azoxystrobin 18.2% w/w + Difenoconazole 11.4% w/w SC is a broad-spectrum systemic fungicide which is recommended for the control of Anthracnose & Powdery Mildew of Chilli; Early Blight & Late Blight of Tomato; Blast & Sheath blight of Paddy; Blight & Downy Mildew of Maize; Rust & Powdery Mildew diseases of Wheat; Leaf spot & Grey mildew of Cotton; Leaf blotch, Leaf spot & Rhizome rot of Turmeric; Purple blotch, Stemphylium blight & Downy mildew of Onion and Red rot, Smut & Rust of Sugarcane.

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For agriculture use.

Re-entry:

Do not enter treated areas for 24 hours after each spray.

Chemical Composition :

Content % w/w

1,2-benzisothiazol-3 (2H)-one	Preservative	0.03 % min.
Water	Solvent	Q.S. % min.
Azoxystrobin a.i.	a.i	18.2 % max.
Cetyl/stearyl-alcohol ethoxilated	Wetter	20.7 % min.
Linear polysaccharide	Thickner	0.25 % min.
Difenoconazole a.i.	a.i	11.4 % max.
Naphthalenesulfonic acid, sodium salt condensated with formaldehyde	Dispersant	1.22 % min.
Blockcopolymer PO/EO	Dispersant	1.13 % min.

Azoxystrobin 18.2% w/w Difenoconazole 11.4% w/w SC

व्यापक शक्ति का अतिरिक्त, यह फंगिसाइड है।

Broad Spectrum Systemic Fungicide

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।



कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

Keep out of reach of children.

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

Keep in cool, dry place away from heat and open flame.

Regn. No. 100/100/100

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

व्यापक शक्ति का अतिरिक्त, यह फंगिसाइड है।

खतरा

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

व्यापक शक्ति का अतिरिक्त, यह फंगिसाइड है।

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

व्यापक शक्ति का अतिरिक्त, यह फंगिसाइड है।

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

खतरा

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

व्यापक शक्ति का अतिरिक्त, यह फंगिसाइड है।

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

व्यापक शक्ति का अतिरिक्त, यह फंगिसाइड है।

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

व्यापक शक्ति का अतिरिक्त, यह फंगिसाइड है।

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

कृषि क्षेत्रों में उपयोग के लिए सुरक्षित है।

Safety Pictograms:



TM	©n TM ©a œ I «	SISV	SIT	XSS	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q	UWS	VX	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q
² ²	m TM m ^{TM2} TM ©a · µ ú © m ^{TM2} TM	SISV	SIT	XSS	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q	UWS	X	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q
—	° »° 2 TM ©a ® tÿ « m ^{TM2}	SISV	SIT	XSS	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q	UWS	VT	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q
³ i	i 2 — ®ñ Ý « TMZ ° µú@µ 3 2 — « ©a 2 « 2 µ — «	SISV	SIT	XSS	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q	UWS	Z	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q
² á—	m ^{TM2} TM ©a 2 « 2 µ — «	SISV	SIT	XSS	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q	UWS	UY	®µ ÝÊ— TM n ° ®µ 3 TM—, n ©— ° 2 3 2 —Q «° ÝÊ— TSZ TX «—m© µ ®° 2 — © ©—n —®— —Q

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Azoxystrobin 18.2% w/w Difenconazole 11.4% w/w SC

(Broad Spectrum Systemic Fungicide)

Azoxystrobin 18.2% w/w + Difenconazole 11.4% w/w SC is a broad-spectrum systemic fungicide which is recommended for the control of Anthracnose & Powdery Mildew of Chilli; Early Blight & Late Blight of Tomato; Blast & Sheath blight of Paddy; Blight & Downy Mildew of Maize; Rust & Powdery Mildew diseases of Wheat; Leaf spot & Grey mildew of Cotton; Leaf blotch, Leaf spot & Rhizome rot of Turmeric; Purple blotch, Stemphylium blight & Downy mildew of Onion and Red rot, Smut & Rust of Sugarcane.

Chemical Composition:	Ingredient	%W/W
1,2-benzisothiazol-3 (2H)-one	Preservative	0.03 % min.
Azoxystrobin a.i.	a.i	18.2 % max.
Blockcopolymer PO/EO	Dispersant	1.13 % min.
Cetyl/stearyl-alcohol ethoxilated	Wetter	20.7 % min.
Difenconazole a.i.	a.i	11.4 % max.
Linear polysaccharide	Thickner	0.25 % min.
Naphthalenesulfonic acid, sodium salt condensated with formaldehyde	Dispersant	1.22 % min.
Polydimethyl siloxane	Anti foaming agent	0.79 % min.
Water	Solvent	Q.S. % min.
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water Litre/ha	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI g a.i./ha	Formulation ml/liter					
Cotton	Leaf spot & Grey mildew	0.03%	0.1%	500	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence.	24 hours	12	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence. Maximum two applications per season.
Sugarcane	Red rot, Smut & Rust	0.03%	0.1%	400	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence.	24 hours	265	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence. Maximum two applications per season.

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

Wheat	Rust & Powdery mildew	0.03%	0.1%	500	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence.	24 hours	35	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence. Maximum two applications per season.
Tomato	Early blight & Late blight	0.03%	0.1%	500	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence.	24 hours	5 days	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence. Maximum two applications per season.
Paddy	Blast & sheath blight	0.03%	0.1%	500	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence.	24 hours	31	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence. Maximum two applications per season.
Onion	Purple blotch, Stemphylium blight & Downy mildew	0.03%	0.1%	500	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence.	24 hours	7	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence. Maximum two applications per season.
Maize	Blight & Downy Mildew	0.03%	0.1%	500	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence.	24 hours	26	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 10 14 days depending upon disease incidence. Maximum two applications per season.

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Chilli	Anthraco nose & Powdery Mildew	0.03%	0.1%	500	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence.	24 hours	5 days	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence. Maximum two applications per season.
Turmeric	Leaf blotch, Leaf spot & Rhizome rot	0.03%	0.1%	500	First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence.	24 hours	60	First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence. Maximum two applications per season.

Note:-

Good Practices relating to pesticide application techniques

Direction of Use

Take the required quantity of fungicide, dilute with a little water and stir using a stick. Then add the remaining quantity of water as per spray volume required and mix well again by stirring.
First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 15 days depending upon disease incidence. Maximum two applications per season.

Weather Condition(in general):

Do not spray during the windy weather conditions or during hot hours of the day, if the foliage is wet or if rain is imminent. Do not spray during adverse crop conditions or when crop is under stress. Spray strictly with the wind to prevent prolonged or repeated exposure to the ~~Soil & Water (where ever applicable)~~ by overlapping applications or by exceeding the recommended rate.

This is not a soil applied fungicide hence, not applicable. Do not contaminate water used for irrigation or domestic purpose and water areas viz. ponds, ditches, lakes, drainage systems, etc, by disposal of product waste, direct over spray or spray drift.

Crop Stage for application:

First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 15 days depending upon disease incidence.

Application Technique

Foliar Spray: Spray with knapsack or any other suitable sprayer.

Equipment Type:

Spray with knapsack or any other suitable sprayer.

Nozzle Type:

Hollow cone nozzle or any other suitable nozzle.

Dosage:

Please refer recommendation table.

Instruction for Mixing:

Please see " direction of use"

Re-entry Period

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

24 hours

Manner and Time of Application

First application to be given as preventive or at disease initiation; repeat one spray at the spray interval of 10 . 14 days depending upon disease incidence.

Application Timing

Should be used preferably in early morning or evening hours

Frequency

First application to be given as preventive or at disease initiation and repeat one spray at the spray interval of 10 . 14 days depending upon disease incidence. Maximum two applications per season.

Stages during which pesticides should not be applied on any crop :

Not to be applied during crop maturity and just before crop harvest

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

Symptoms Of Poisoning

Poisoning may cause symptoms of activity decrease, Pupil dilation, moderate foot withdrawal, reduced splay reflex, increased breathing depth and reduced breath rate. etc.

First Aid

Remove the affected person to a well-ventilated room or to fresh air and protect him from under cooling. In case of suspected poisoning, stop work immediately and call a physician.

If ingested: if person is conscious and alert, rinse mouth with water and give 1 or 2 glass of water to drink. Never give anything by mouth to an unconscious person. Call the doctor. Do not induced vomiting until and unless desired by attending physician.

If inhaled: Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration,Consult a physician.

In case of skin contact: Wash thoroughly the affected part of body with soap and plenty of water including hairs and under fingernails. If pain, redness or irritation persists, consult a doctor. All contaminated clothes should be segregated and thoroughly washed with soap and water separately before use.

In case of eye contact: Rinse gently with plenty of water at least for 15 minutes. If pain, irritation, redness or photophobia persists. Consult eye specialist

Antidote

No specific antidote is known; apply symptomatic therapy.

Phytotoxicity

Not phytotoxic at recommended dosage. recommendation

Storage Conditions

1. The packages containing the fungicide should be tightly closed, stored in original containers in separate rooms or premises away from the rooms or premises used for storing other articles particularly articles of food or animal feed or shall be kept in separate almirahs under lock and key, away from children, unauthorized persons and animals.

2. The rooms or premises meant for storing the fungicide shall be well built, dry, well lit and ventilated and of sufficient dimension to avoid contamination with vapor. Keep away from direct sunlight.

3. Do not allow the product to freeze.

Safety Precaution:

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

Do not use cooking utensils for preparing the spray solution. Use wooden stick for stirring the spray solution. Avoid contact with skin, eyes, mouth and clothing. Avoid inhalation of fog, vapors or spray mist. Wear chemical resistant hand gloves to avoid contact with the skin. Wear protective clothing (full sleeved shirt and pants), mask, goggles, cap/hat and boots, while handling the product. Do not eat, drink or smoke while applying the product. Wash hands with soap and plenty of water and change clothes after the work is over. Avoid contamination of air and water bodies with the fungicide. Also wash the contaminated clothes before re-use. Newly treated areas must not be grazed and all livestock's area is to be kept out until the treated area is dry. This product should not be applied (tank mix) with any other chemical. Do not blow nozzle with mouth. Do not spray crops under stress. If any indisposition occurs, stop the work, follow first aid measures and call a physician immediately. Spilt product must not be re-used. Collect up spilt material thoroughly and bury it in a safe place. Rinse spilt area with plenty of water.

Cautionary Statement:

- 1. Not to be used for any purpose other than those subscribed on labels & leaflets.**
- 2. The product shall not be used in area where pisciculture/intercropping pisciculture is practiced along with rice cultivation.**
- 3. The combination product is toxic to fish and aquatic organisms therefore, not to be used near aquaculture.**
- 4. Destroy the container after use as directed on leaflet.**
- 5. Toxic to earthworms.**

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufactures, formulators of insecticides and operators to dispose packages or surplus material and washings in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

Plot No. 1, Swastik Ceramic Compound, AT & PO Sankarda,
District Baroda, Gujarat
Maharashtra

Syngenta India Limited, Santa Monica Works, Corlim, Ilhas, Goa
Maharashtra

Unit II, Plot No. 155/1, GIDC Estate, Ankaleshwar, District-
Bharuch, Gujarat
Maharashtra

Manufactured by

Syngenta India Limited
Syngenta India Limited, Amar Paradigm, S.No. 110/11/3, Baner
Road, Pune- 411 045 Pune Maharashtra

Customer Care Details:

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "