

WP(Strain : BB- ICAR - RJP), MCC-1022

1 10 gm min

WP(Strain : BB- ICAR - RJP), MCC-1022

WP(Strain : BB- ICAR - RJP), MCC-1022	1 x 10 <sup>8</sup> / 2 <sup>3</sup> -!	1.15 % wp
WP(Strain : BB- ICAR - RJP), MCC-1022		90.85 % w/w
WP(Strain : BB- ICAR - RJP), MCC-1022		8 % w/w
TSS!SSS~ ± ± ±		

μ	μ	â³ 2		μ	μ	μ	n
		%	%				
		! ñ</td <td>œ</td> <td>*(#ž+(#μ</td> <td></td> <td>\$%š</td> <td></td>	œ	*(#ž+(#μ		\$%š	

Note:-

q™ μW «

1.15% |ø³ @ p' wœ p' @. 3 2 ° @ μ 3 2 , -ž ° -œ 3 ž @ 3 ž -ú , -ž -@

2 ° 2 ° a ©

m-μ 2 ° 2 ° a © 2 q™ -3 i -œ »W | mñ 2 ° 2 ° a © 2 ° -q™ -» -3 i -œ »W

2 œ | ©

»y-° ± 2 œ | ©

° μ m a

n · « - ©

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

2.5 kg in 750-850 lit

\_\_\_\_\_

-

\_\_\_\_\_

\$%\$α

—α p° '®μ' —'m©' μ

-

\_\_\_\_\_

-

n . « —' —' 2 3

° ° » | ' ' , 2 ' —' 2 3

\_\_\_\_\_

-

\_\_\_\_\_

-

\_\_\_\_\_

-

. 1' —' μ ¼'

° ' « « Ž 2 ©μ Ž qýα Ž j μ —' | ' ' wμ j ' ' » ' ° — © ' » Q

® « ®' n . 1 μ ©

-

\_\_\_\_\_:

\$! m™' —α—, —' —™μ' μ³ ' » ' © '™μ' —'®Ÿ'™«™«' —' qýα' —'. wQ%³ «' ñ œ' | ' ' —®E' «' 1 ©' » '™W' »' ©' «' 1 ©' ñ œ' | ' ' —®E' —'®—' | ' ' ° ° —' ° ° —' wQ&!³ «' n —' «' 1 ©' » '™B' ' »' ©' n —' —' ° 2 °³ ' —2 —' | ' '®—' ° ° \$# ' ° ° \$( ' 2 —α' —' μw —' wQ' ! '™' —' © j ' »' . ' 2 ' μ' j ' 3 Q

\_\_\_\_\_

μ ¼' —' q®œ' —'

° » —' —' ©

1' —α—, —' |ø' —' mμ™z' mμ™ —' 2 ' 3 ' n μ² ' 3 ' ° ° j —' 2 ' mō³'. ©w . , 1 —' —' 3 '®«' a | 3 ' ®' n » ' ° ° » ©' —w j ' ©' » ' ° ° «' mō³' —' 2 ' ° ° «' mμ™z' mμ™ —' 2 ' 2 ' 3 ' n μ² ' 3 ' 2 ' ° ° μ' œ° ' 2 ' ° ° «'™ Q2' j —' 2 ' m³ . ' n μ² ' 3 ' 2 ' —α—, —' —' ± | ' ' —' —' »' . »' mçŸ' ©' »' ° ° —' » n Ž ° ° Ž Ÿ | Ž —, ' 3 a © Ž . «' Ž . '®³ b ©' μ ú° ' œ | ' ' » —' œ » 3 ' © —' —α—, —' —' ± ®' ° ° . © . '™' «' 1™' » —' —' | ' ' —' » Q

\_\_\_\_\_:

Caution: **Not to Be Used on Crops Other Than Specified on This Label/Leaflet**



**Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022**

**( Contact Insecticide )**

**1 10 gm min**

Beauveria bassiana is based on Etamopathogenic fungi containing Beauveria bassiana-1.15% which controls Rice leaf folder in rice.

<b>Chemical Composition:</b>	<b>Ingredient</b>	<b>%W/W</b>
Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022	1X10 <sup>8</sup> / gm min	1.15 % wp
Carrier (talc)		90.85 % w/w
Moiture		8 % w/w
<b>Total:</b>		<b>100.000% w/w</b>

**Recommendation**

<b>Crop(s)</b>	<b>Common Name of Pest</b>	<b>Dosage/HA</b>		<b>Dilution in Water Liter</b>	<b>Crop Stage</b>	<b>Re-entry Period</b>	<b>Pre Harvest</b>	<b>Frequency</b>
		<b>AI %</b>	<b>Formulation kg</b>					
rice	Leaf folder (Cnaphalocrosis medinalis)	1.15%WP	2.5kg	750-850 liter	-	12 hours	-	-

**Note:-** -

**Good Practices relating to pesticide application techniques**

**Direction of Use**

Beauveria bassiana-1.15% WP can be sprayed in the field by using the equipment like backpack hollow type sprayers machine, bucket sprayers, pump sprayers, compression, knapsack battery sprayers at ASPEE HPT power sprayers.

**Weather Condition(in general):**

should be used in favorable weather condition and should not be used in extreme weather conditions

**Soil & Water(where ever applicable):**

Light to heavy soil and clean water

**Crop Stage for application:**

-

**Application Technique**

-

**Equipment Type:**

=

**NozzleType:**

=

**Dosage:**

2.5 kg in 750-850 lit

**Instruction for Mixing:**

-

**Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet**

**Re-entry Period**

12 hours

**Pre-Harvesting Interval**

-

**Manner and Time of Application**

-

**Application Timing**

morning and evening hours.

**Frequency**

-

**Stages during which pesticides should not be applied on any crop :**

=

**Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :**

-

**Symptoms Of Poisoning**

Headache, Nausea, Vomiting, irritation & allergy may occur.

**First Aid**

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to 15 minutes. 4. Remove the patient into fresh air.

**Antidote**

Treat symptomatically.

**Phytotoxicity**

-

**Storage Conditions**

1. The packages containing the insecticides shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles or shall be kept in separate almira under lock and key depending upon the quantity and nature of the insecticide. 2. The rooms or premises meant for storing the insecticides shall be well built, dry, well-lit, ventilated and of sufficient dimensions to avoid contamination with vapours.

**Safety Precaution:**

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothing which includes rubber gloves, boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

**Cautionary Statement:**

**1. Not to be used for any purpose other than those subscribed on label and leaflet. 2. Destroy the container after use as directed on leaflet.**

**Disposal of used containers, washing from equipment and surplus material :**

**Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet**

1 It shall be the duty of manufacturers, formulators of Insecticides and operators to dispose packages or surplus materials and washings in a safe manner so as to prevent Environmental or water pollution. 2 The used packages shall not be left outside to prevent their re-use. 3 Packages shall be broken and buried away from habitation.

**Manufacturer Premises :**

Gut No. 280 Budhgaon Bisur Road Sangli 416304 Maharashtra  
Maharashtra

**Manufactured by**

Sarthak Agro Laboratories  
H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Maharashtra

**Customer Care Details:**

Sarthak Agro Laboratories  
H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Sangli Maharashtra

**Safety Pictograms:**



**Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "**

**Primary Package Label**

**[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]**

**Read leaflet before use:**

**Common Name:**

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

**BRAND NAME: -**

**Chemical Composition :**

**Content % w/w**

Beauveria bassiana 1X10<sup>8</sup> / gm min 1.15 % wp  
1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Carrier (talc) 90.85 % w/w

Moiture 8 % w/w

**Total: 100.00%**

**Please refer leaflet for details**

**Antidote**

Treat Symptomatically.

**Manufactured by**

Sarthak Agro Laboratories  
H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Sangli Maharashtra

**Premises Address**

(1). Gut No. 280 Budhgaon Bisur Road Sangli 416304 Maharashtra

**Customer Care Details**

Sarthak Agro Laboratories  
H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Sangli Maharashtra

**Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022**

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

**Contact Insecticide**

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022



Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Keep in cool, dry place away from heat and open flame.

Regn. No. \_\_\_\_\_

Mfg. Lic. No. \_\_\_\_\_

Batch No. \_\_\_\_\_

Mfg Date \_\_\_\_\_

Exp. Date \_\_\_\_\_

Net Content / Volume \_\_\_\_\_

Max. Retail Price \_\_\_\_\_

Customercare \_\_\_\_\_

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

**Safety Pictograms:**



**Secondary Package Label**

**[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]**

**Read leaflet before use:**

**Common Name:**

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

**BRAND NAME: -**

**Crops and Pest Groups: Rice-Leaf folder caphalocrosis medinalis**

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

**Purpose:**

for agricultural use

**Re-entry:**

12 hours

**Chemical Composition :**

**Content % w/w**

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022	1X10 <sup>8</sup> / gm min	1.15 % wp
Carrier (talc)		90.85 % w/w
Moiture		8 % w/w

**Total: 100.00%**

Safety Precautions, Symptom of Poisoning and First Aid : Please refer leaflet for details.

**Antidote Statement:**

Treat Symptomatically.

**Cautionary statement:**

- Not to be used for any purpose other than those subscribed on label and leaflet
- Destroy the container after use as directed on leaflet.

**Manufactured by**

**Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022**

Beauveria bassiana 1.15% WP, (Strain : BB- ICAR - RJP), MCC-1022

**Contact Insecticide**

300g



Keep in cool, dry place away from heat and open flame.

Keep in cool, dry place away from heat and open flame.

Regn. No. \_\_\_\_\_

Mfg. Lic. No. \_\_\_\_\_

Batch No. \_\_\_\_\_

Mfg Date \_\_\_\_\_

Exp. Date \_\_\_\_\_

Net Content / Volume \_\_\_\_\_

Max. Retail Price \_\_\_\_\_

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

300g

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Contact Insecticide

300g

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

300g

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

300g



**Product Address**  
Sarthak Agro Laboratories  
H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Sangli Maharashtra

(1).

»—« ± μ . . . . .

° ° ± w . . . . . μ ° . . . . .  
H.No. 3351Z -3 . . . . . μ → 03 . . . . . μ · È . . . . . © . . . . . μ γ μ · Z · © . . . . . Z · ý . . . . .  
. . . . . μ . . . . . 2 »

**Customer Care Details**

Sarthak Agro Laboratories  
H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Sangli Maharashtra

**Safety Pictograms:**



## Big Label

(> 250 mL Primary and Secondary Package Label)

**Read leaflet before use:**

**Common Name:**

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

**BRAND NAME: -**

**Crops and Pest Groups: Rice-Leaf folder caphalocrosis medinalis**

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

**Purpose:**

for agricultural use

**Re-entry:**

12 hours

**Chemical Composition :**

**Content % w/w**

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022	1X10 <sup>8</sup> / gm min	1.15 % wp
Carrier (talc)		90.85 % w/w
Moiture		8 % w/w

**Total:100.00%**

**Safety Precautions:**

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothing which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

**Symptom of Poisoning:**

**Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022**

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Contact Insecticide

सावधान



Keep in cool, dry place away from heat and open flame.

Regn. No.		
Mfg. Lic. No.		
Batch No.		
Mfg Date		
Exp. Date		
Net Content / Volume		
Max. Retail Price		

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

1.15% WP

90.85% w/w

1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

90.85 % w/w

8 % w/w

TSSISS

Beauveria bassiana 1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

1.15% WP (Strain : BB- ICAR - RJP), MCC-1022

90.85% w/w

8% w/w

TSSISS

Headache, Nausea, Vomiting, irritation & allergy may occur.

**First Aid:**

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to 15 minutes. 4. Remove the patient into fresh air.

**Antidote**

Treat Symptomatically.

**Cautionary statements:**

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

**Manufactured by**

Sarthak Agro Laboratories  
H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Sangli Maharashtra

**Premises Address**

(1). Gut No. 280 Budhgaon Bisur Road Sangli 416304 Maharashtra

**Customer Care Details**

Sarthak Agro Laboratories  
H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Sangli Maharashtra

**Safety Pictograms:**



© 2018 Sarthak Agro Laboratories

Warranty: The product is warranted to be free from defects in material and workmanship under normal use and conditions. The warranty is void if the product is used for any purpose other than that specified on the label and leaflet. The warranty is also void if the product is used in any manner which is not recommended by the manufacturer.

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Warranty: The product is warranted to be free from defects in material and workmanship under normal use and conditions. The warranty is void if the product is used for any purpose other than that specified on the label and leaflet. The warranty is also void if the product is used in any manner which is not recommended by the manufacturer.

© 2018 Sarthak Agro Laboratories

© 2018 Sarthak Agro Laboratories

(1).

Warranty: The product is warranted to be free from defects in material and workmanship under normal use and conditions. The warranty is void if the product is used for any purpose other than that specified on the label and leaflet. The warranty is also void if the product is used in any manner which is not recommended by the manufacturer.

H.No. 3351, Near Lokmanya Balwadi Thote Galli, Tal - Walwa  
Walwa Sangli Maharashtra