

313001

Premises Address

- (1). 237, GIDC Industrial Estate, Panoli, District Bharuch, 394116, Gujarat
- (2). Plot No 3133 to 3139 and others, GIDC Estate, Panoli, District Bharuch, Gujarat
- (3). Udaisagar Road, Udaipur-313001, Rajasthan

© 2014 P I Industries Ltd. All rights reserved.

- (1).
- (2).
- (3).

Customer Care Details

Address: P I Industries Ltd, Udaisagar Road, Udaipur-313 001,
Contact no: 0124-6780290
Email: sm.ahsan@piind.com

» « ± μ · · · »

Address: P I Industries Ltd, Udaisagar Road, Udaipur-313 001,
Contact no: 0124-6780290
Email: sm.ahsan@piind.com

Safety Pictograms:



2. «-|' @-

3. »'»°° o|'®' ln·«-— μw»'| 'm-, ° @- '—®' q®3™ →w| -'®'o° —p' ©—μ' ±·'—»' »Qj μ³—1'—'®°' q®3™ —'—'°° œ

-° μ'm- a

-° l' n³ '®' - \$z\$% DAS

n·«- '©— —

®' ln·«- qœ· ýπj ' 3' | 'vw° @pžwœ®'®· ' 3' —' q®3™ —'—' μπ'®' 3' ° È'j π'—j μ'—'° a' μ™3' j' ©' »Q

÷μπ'—'—' μj' Åπ'—j μ'—'° a' —'—' 3' Q

÷μπ'—'—' μj' Åπ'—j μ

Rice (Nursery/ transplanted/ Direct seeded) :
200-250ml formulation /ha.(20-25 ga.i./ha.) in 300lit. of water

q®3™ —'—'® μ'° ©μ'—' »μ vQ»°° o|'—' n· 3—2 —'—2 ®| 'o° —2'2 —'2'®—'2'2 μ vQ q®3 a'®' 3' —'° a'—'° μ' —'—' μ'—'—' μw®'®' »'° n « μ'—'° a'—' «l π'®'—'—', 1'2 —' j' ÈQ

% 'š π

®- , μ'™ ®®- —'—' μw' q œ°° 2 3' »' j' °' m- —'—' ®®· ' q™—'—'® œ° mª ®-° l' —'—' μw TSzTU' «- '° n p'—'® œ®Z w' '®®- —'—' μw TSzTW «- '® p'—'® œ®Z ÝÈ—'—' Q m®' V'° W®' —' m· a' 2' q®3™ —' Q 3' »° 2 3' 2 π' w' . ©· ' 3' ®' a' ®3' —' m- ®' ÝÈ—'—' Q

°- '° j' p'· μ'—'—' μw' ÝÈ—'—' q œ°° 2 3' ° n p'—' TXZUX' «- '® œ®' »Q

n·«- —'° 2 3

m-2 -©'°° »| ' , 2' —'° 2 3 Q

w—'° ' n·«-

n·«- —'° 2 3 —' q®3 a'®' -« , —'® μ'—'—'

3 »' qñ®«' 2 Ý μ³ —' μw' . 1 á®' »| ' 'o° —'° ©2 μ' wá· —'ýœ'—' n °®°'—' »—' 3' j' —'œ »vQ

. 1'—' μ¼' '

—2 j' ž qýπ ž 2 »'° μ' n —'—' m° ' Q

®«®' n . 1 μ©

°- ' , —' m-° ' q®3™ →w| -'®' 3 »' qñ®«' ' oπ π á° —'—' »' »Q

Caution:Í Not to Be Used on Crops Other Than Specified on This Label/LeafletÍ "



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Bispyribac Sodium 10% SC**(Systemic Herbicide)**

Bispyribac Sodium 10% SC is a post emergence broad spectrum herbicide effective against grasses, sedges and broad leaf weeds of rice crop.

Chemical Composition:	Ingredient	%W/W
Aluminium Oxide C		1.50 % w/v
Bispyribac Sodium	active ingredient	10.00 % w/v
Polyalkylene oxide derivative of a synthetic alcohol	Adjuvant A	31.90 % w/v
Polyxyethylene polyoxy Propylene Synthetic alcohol ether	Adjuvant B	53.20 % w/v
Propylene glycol		2.00 % w/v
Silicon KS-506		1.00 % w/v
Stabilizer		1.00 % w/v
Water		to make 100 QS
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water Litre	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI gms.	Formulation ml.					
Rice(Direct Seeded)	Fimbristylis miliaceae Eclipta albe Ludwigia parviflora Monochoria vaginalis Alternenthere phyloxiroides Sphenoclea zeylanica	20-25gm	200-250ml	300 Liter	10-15 DAS	24 hr.	78 days	one spray
Rice (Transplanted)	Ischemum rugosum Cyperus difformis Cyperus iria	20-25	200-250	300 Liters	10-14 DAP	24 hr.	78 days	one spray
Rice (Nursery)	Echinochloa crusgalli, Echinochloa colonum,	20	200	300	Nursery - 10-12 DAS	24 hr	-	one time application

Note:- .**Good Practices relating to pesticide application techniques****Direction of Use**

use recommended dose. Dilute the formulation in required quantity of water

Weather Condition(in general):

Do not spray during rainy, windy and hot sunny period

Soil & Water(when ever applicable):

This Herbicide is for foliar application and has no adverse effect when used at recommended dose . Avoid using near aquaculture

Crop Stage for application:

Nursery - 10-12 DAS

Application Technique

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

Foliar application using high volum sprayers and ASPEE-HTP power sprayer fitted with flat fan or flood jet nozzle.

Equipment Type:

knapsack sprayer with flat fan/ flood jet nozzle.

NozzleType:

flat fan/ flood jet nozzle.

Dosage:

Rice (Nursery/ transplanted/ Direct seeded) :
200-250ml formulation /ha.(20-25 ga.i./ha.) in 300lit. of water

Instruction for Mixing:

Shake the bottle before use measure out requisite quantity of the herbicide and mix it well with small quantity of water. Add the remaining quantity of water as specified with thorough agitation for total coverage of crop with suitable sprayer.

Re-entry Period

24 hr

Manner and Time of Application

Nursery/ Transplanted Rice: The optimum time of application is 10-12 days after sowing for nursery rice and 10-14 days after planting for transplanted rice when most of the weeds have already emerged and are of 3- 4 leaf stage depending upon soil and climatic factors.

Direct seeded Rice: For direct seeded rice optimum time of application is 15-25 days after sowing.

Application Timing

Preferably in the morning and evening hours.

Frequency

one time application

Stages during which pesticides should not be applied on any crop :

Follow the above instructions as mentioned in time of application

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

This product is toxic to fish and not be used around aquaculture.

Symptoms Of Poisoning

Hypo-activity, vomiting, salivation and diarrhoea.

First Aid

In case of Inhalation: Immediately move victim to fresh air. If breathing is irregular or stopped, administer artificial respiration. Keep patient warm and at rest. Call doctor immediately.

In case of Ingestion: If swallowed, seek medical advice immediately. Do not induce vomiting. Never give anything by mouth to an unconscious person.

In case of skin contact: Take off all contaminated clothing immediately and thoroughly wash the affected parts of the body with plenty of water. Wash contaminated clothes before re-use.

In case of eye contact: Rinse immediately with plenty of water, also under the eyelids for atleast 15 minutes. If pain or irritation persists, consult eye specialist.

Antidote

No specific antidote is known. Treat symptomatically

Phytotoxicity

The product is not phototoxic when used as per field recommendation.

Storage Conditions

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

1. The package containing the herbicide shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles particularly articles of food or shall be kept in separate almirahs under lock and key depending upon the quantity and nature of the herbicide.
2. The rooms or premises meant for storing the herbicide should be well built dry, well . Ait and ventilated and of sufficient dimension to avoid contamination by vapour.

Safety Precaution:

1. Keep away from foodstuffs, empty foodstuff containers and animal feed.
2. Avoid contact with mouth, eyes and skin.
3. Avoid inhalation of the spray mist.
4. Wash thoroughly the contaminated clothes and parts of the body after spraying.
5. Do not smoke, drink, eat and chew anything while spraying.
6. Use protective clothing which includes long sleeved shirt and long pants, chemical resistant gloves, shoes, goggles and socks while spraying.

Cautionary Statement:

-Not to be used on crops other than those prescribed on label and leaflets.

-Destroy the container after use as directed in the leaflet.

-This product is toxic to fish and not be used around aquaculture.

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufactures, formulators of insecticides and operators to dispose packages or surplus material and washings in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

237, GIDC Industrial Estate, Panoli, District Bharuch, 394116,
Gujarat
Haryana

237, GIDC Industrial Estate, Panoli, District Bharuch, 394116,
Gujarat
Rajasthan

Plot No 3133 to 3139 and others, GIDC Estate, Panoli, District
Bharuch, Gujarat
Haryana

Plot No 3133 to 3139 and others, GIDC Estate, Panoli, District
Bharuch, Gujarat
Rajasthan

Udaisagar Road, Udaipur-313001, Rajasthan
Haryana

Udaisagar Road, Udaipur-313001, Rajasthan
Rajasthan

Imported and Repacked by

P I Industries Ltd
4th Floor, Tower-A, Mellennium Plaza, Sector-27, Gurgaon-122 002
Gurgaon Haryana

Customer Care Details:

Address: P I Industries Ltd, Udaisagar Road, Udaipur-313 001,
Contact no: 0124-6780290
Email: sm.ahsan@piind.com

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "