

Primary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Fenpyroximate 5% SC

BRAND NAME: -

Chemical Composition :

		Content % w/w
Thickener (Xanthum Gum) (Heteropolysaccharide)	Thickener	4.0 % w/w
Anti-freezing agent (Propylene glycol)	Anti-freezing agent	5.0 % w/w
Fenpyroximate a.i. content	active ingredient	5.0 % w/w
Defoamer (Dimethyl polysiloxane)	Defoamer	0.25 % w/w
Dispersing agent (Polyoxyethylene styryl phenyl ether and polyoxyethylene alkyl aryl ether)	Dispersing agent	6.0 % w/w
Preservative (1,2-benzisothiazin-3-one)	Preservative	0.1 % w/w
Distilled water	other ingredient	Q.S. to make 100% w/w
Total:		100.00%

Please refer leaflet for details

Antidote

there is no specific antidote. hence, symptomatic and supportive therapy should be given.

Manufactured by

Fenpyroximate 5% SC

फेनपिरोक्सिमेट 5% SC

Acaricide

अकारिड



Keep out of reach of children.

Keep in cool, dry place away from heat and open flame.

Keep in cool, dry place away from heat and open flame.

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Net Content / Volume _____

Max. Retail Price _____

Customercare _____

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

फेनपिरोक्सिमेट 5% SC

Secondary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Fenpyroximate 5% SC

BRAND NAME: -

Crops and Pest Groups: -

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

for agricultural use.

Re-entry:

48 hours after spraying.

Chemical Composition :

Content % w/w

Thickener (Xanthum Gum) (Heteropolysaccharide)	Thickener	4.0 % w/w
Anti-freezing agent (Propylene glycol)	Anti-freezing agent	5.0 % w/w
Fenpyroximate a.i. content	active ingredient	5.0 % w/w
Defoamer (Dimethyl polysiloxane)	Defoamer	0.25 % w/w
Dispersing agent (Polyoxyethylene styryl phenyl ether and polyoxyethylene alkyl aryl ether)	Dispersing agent	6.0 % w/w
Preservative (1,2-benzisothiazin-3-one)	Preservative	0.1 % w/w
Distilled water	other ingredient	Q.S. to make 100% % w/w

Total: 100.00%

Fenpyroximate 5% SC

फेनपिरोक्सीमेट ५% एससी

Acaricide

खतरा



बच्चों के दूरी में रखें।

Keep out of reach of children.

ठंडी, सूखी जगह पर रखें।

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Net Content / Volume _____

फेनपिरोक्सीमेट ५% एससी

फेनपिरोक्सीमेट ५% एससी

खतरा

खतरा

फेनपिरोक्सीमेट ५% एससी

फेनपिरोक्सीमेट ५% एससी

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

खतरा

फेनपिरोक्सीमेट ५% एससी

Safety Precautions, Symptom of Poisoning and First Aid :Please refer leaflet for details.

Antidote Statement:

there is no specific antidote. hence, symptomatic and supportive therapy should be given.

Cautionary statement:

- 1. not to be used for any purpose other than subscribed on label & leaflets.
2. destroy container after use as directed on leaflet.
3. do not use in those situations if there is possibility of harming bees & fish, livestock, game, wild birds & animals.

Manufactured by

Rallis India Limited
Rallis India Limited, IIIrd Block, IIInd Floor, KSCMF Ltd., Building, 8, Cunningham Road, Bangalore - 560052 Rallis India Limited 156/157, Nariman Bhavan, 15th Floor, 228, Nariman Point, Mumbai - 400021 Bangalore Bangalore Karnataka 560052

Premises Address

- (1). M/s Rallis India Limited, Plot No. 3301, GIDC Industrial Area, Dist. Bharuch, Ankleshwar-393 002, GUJARAT Gujarat
(2). M/s Rallis India Limited, Plot No. 2808, G.I.D.C. Estate, Ankleshwar - 393002, Dist. Bharuch, GUJARAT Gujarat
(3). M/s Rallis India Limited, C-5/C-6, MIDC Industrial Area,Phase-III, Shivani, Akola . 444 104, MAHARASTHRA Maharashtra
(4). Rallis India Ltd Plot No. C44, Chemical Zone, Dahej Taluka-Vagra, District- Bharuch - 392130, GUJARAT

Customer Care Details

Address: Rallis India Limited, IIIrd Block, IIInd Floor, KSCMF Ltd., Building, 8, Cunningham Road, Bangalore - 560052, Email: kjsrivalli@rallis.co.in, Contact No: 09810991089

Max. Retail Price m - 02 - 2 y3

qñ®«—

µ ° o ð³ µ ² µ
µ ° o ð³ µ ² µ Ž 3 µ — Ž 2 ÷ µ —!w! °!w! ! µ! µ! õ™Ž
8Ž —õ—õ™»² µ Ž ° õ™µ . 560052 µ ° o ð³ µ ² µ Ž 156/157Ž
— ² — ± . — Ž 15 ÷ µ Ž 227 — ² — ° õõŽ ² ú° p! . 400021 ° õ™µ
560052

@ ° — — @ _

(1). ????? ????????? ???????, ????? ?? 3301, ???.??.??.??.
???????????????? ??????, ????? ?????, ????????????? 393002, ???????

(2). ????? ????????? ???????, ????? ?? 2808, ???.??.??.??.
???????????????? ??????, ????? ?????, ????????????? 393002, ???????

(3). ????? ????????? ???????, ??-5/??-6, ???.??.??.??. ???-3, ?????,
????? - 444104, ?????????????

(4). ????? ????????? ???????, ????? ?? .??44, ?????? ???, ?????,
??????-?????, ????? ????? - 392130, ???????

»—« ± µ

@ _ Rallis India Limited, IIIrd Block, IIInd Floor, KSCMF Ltd., Building, 8, Cunningham Road, Bangalore - 560052, p² µ - kjsrivalli@rallis.co.in, Contact No: 09810991089

Safety Pictograms:



Big Label

(> 250 mL Primary and Secondary Package Label)

Read leaflet before use:

Common Name:

Fenpyroximate 5% SC

BRAND NAME: -

Crops and Pest Groups: -

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

for agricultural use.

Re-entry:

48 hours after spraying.

Chemical Composition :

		Content % w/w
Thickener (Xanthum Gum) (Heteropolysaccharide)	Thickener	4.0 % w/w
Anti-freezing agent (Propylene glycol)	Anti-freezing agent	5.0 % w/w
Fenpyroximate a.i. content	active ingredient	5.0 % w/w
Defoamer (Dimethyl polysiloxane)	Defoamer	0.25 % w/w
Dispersing agent (Polyoxyethylene styryl phenyl ether and polyoxyethylene alkyl aryl ether)	Dispersing agent	6.0 % w/w
Preservative (1,2-benzisothiazin-3-one)	Preservative	0.1 % w/w
Distilled water	other ingredient	Q.S. to make 100% % w/w

Fenpyroximate 5% SC

फेनपिरोक्सिमेट 5% SC

Acaricide

खतरा



Keep out of reach of children.

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Net Content / Volume _____

Max. Retail Price _____

फेनपिरोक्सिमेट 5% SC

खतरा

फेनपिरोक्सिमेट 5% SC

Acaricide

खतरा

Keep out of reach of children.

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Net Content / Volume _____

Max. Retail Price _____

फेनपिरोक्सिमेट 5% SC

खतरा

फेनपिरोक्सिमेट 5% SC

Acaricide

खतरा

Keep out of reach of children.

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Safety Pictograms:



—@3'â° 2 ² X' : . µú° —'° ð "

fi' 2 | —' fl

—@3'â° 2 ² X' : . µú° —'° ð " w—m ²@« —, —'»' j ° —' 3 TM 2 œ —' @µ' —, œ³ —' µ µŽ TMµ ° | '° TM —' —' —' a 2' —' µw —³ j ©' »' Q
 ° 3 —'° 'œ—'

m ð² ¢' w' ² fi' @o µ—' äµ—µ fl	m ð² ¢' w' ²	XIS' —' w/w
j' —' 2' fi' 2 a' oµ' @µ' ° µ ä° —' fl	j' —' 2'	SIUX' —' w/w
j' @° TM w' ² fi' µ n' ä° o a' µ—' ² oµ'	j' @° TM w' ²	YIS' —' w/w
—oµ' p a' ' w' ² fi' µ n' ä° o a' µ—' n µ—oµ'		
w' oµ' p a' ' fl		
n ° ©' j µ	m ð³ 'š ²—	â³!w' TSS' —' —' —' w/w
—@3'â° 2 ² X' : . 3' © .	° 3' š ²—	XIS' —' w/w
° ' ². ². fi' Ź j' ° a 3 j' —ž ž —fl	° ' ². ².	SIT' —' w/w
a —' j' ð a 2' TM fl fi' @ µ° —' o j' fl	a —'	WS' —' w/w

3 TM - TSSISS' ± "±'

—° µ'	—² —' —' 2	©' »â³' 2		® —' —' 2 ® —' —' 2' fi' µ' fl	—° µ' m' a	@ —' —' µ	® l' —° µ	n .
		° j' ©ñ ° 3' ©ñ fi' 2 fl	° 'œ—' ° 'œ—' fi' 2' µ fl					
œ³	µ µŽ TMµ ° '° TM —' —'	VSzYS	YSSzTUSS	VßS	j' —' m' a' Q	3 TM —' w' ² š ² —' ° «' »' —' —' Q	Z	—² —' «° . . . n —' ®' UZV' ® Ÿ j' —' —' Q
2 œ	@ µ' —'	TXzVS	VSSzYSS	XSSzZXS	j' —' m' a' Q	3 TM —' w' ² š ² —' ° «' »' —' —' Q	V	—² —' «° . . . n —' ®' UZV' ® Ÿ j' —' —' Q

Note:-

q®³ TM —' µw —' «' .

®« a | —' —' | ©' 2' —' q®³ TM —' Q @ —' —' n . 3 —' 2' j' —' —' | ©' —' TMß »' q° 2' ®« a | mçŸ' —' —' 2' µ w Q

2 ° 2' —' a ©

° ° ©ž »' | '° TM l —' ®' m' —' —' «' —' Ÿ j' —' . " 2 «' ± TM —' —' Q

2 ² | ' © —

3 »' ®« a | ®' Ÿ j' —' —' µw q®³ TM —' ³ j' ©' »' Q o° µw 2' ² —' o° —' ©' —' —' p' —' ©' —' »' »' Q j' µ³ —' 1' 2' | '° —' n ° —' ®°
 q®³ TM —' —' Q

—° µ' m' a

j' —' m' a' Q

Caution: í Not to Be Used on Crops Other Than Specified on This Label/Leaflet "'

n . « - ' © — —

qçœ . ý³² . μ . ³º | — j ' w @ ' ' Ÿ | — . ' — ' Q

- @ - @ ³ ' ž - « ' ³ ' ž — ú , — ' - @ - ' ³ ' ž — ú , — ' - @ - ' « ' ³ ' ž ' ASPEE - HTP ³ ' Q

--

15-30 g a.i./ha (300-600 mL formulation/ha) in 500-750 litres of water on Chilli and 30-60 g a.i./ha (600-1200 mL formulation/ha) in 400 litres of water on Tea.

³ ' TM — W ' š « ' » ' . ' — ' Q

n . « - ' — ' ² ³

º » | ' , ² ' — ² ³ ' o @² μ ' — ' — ' é³ « ' » » © ' » ' Q

— « ' — « ° . ' — n — ' @ ' UZV ' @ ' ' Ÿ | — . ' — ' Q

-

. 1 ' — μ ¼ ' ' .

³ » μ ¼ ' ' o a ' ² ' » ' o — © ' » ' i ' o œ — ž ' « « ž š ' » » ž ' œ — ž ' a — TM © ž μ ' « @ — ž @ « a ' x œ — ž ñ ñ œ ' mº ' ' i ' o ' ' j ' μ ž œ © ž ' a ' É ' o ' ñ ñ œ ' ñ — ' ©² ' ' ' m³ — ' — ' œ — ' » ' o — © ' » ' Q

@ « @ ' n . 1 μ ©

- @ ³ ' âº ² « X ' . μ úº — ' o ð ' — ' o © ³ ' TM ² ' — ' — ' ³ ' TM — ' — ' @ ' — ' μ ' @ ' — ' » ' o ' mº ' — ' » ' @ É © ' » ' Q

T! ' — TM μ ' j ' — ' @ ' — ' ³ ' « ý³ â © » , ² ' | ' — ' — ' © — ž ' © ' @ — ' — ' o a ' ² ' — ' — ' y μ ' — ' | ' qº ' T ' ³ ' U ' TM μº ' @ — ' @ μ w Qº » , ý³ â © — ' ² »º — ' Ÿ ' ± — ' « ' Q | â « ' — ' — ' μ w Q j ' — ' — ' œ — © — ' y « ' — ' — ' » — ' j ' w q y « ' — ' — ' w Q U ' — ' o ' ² ' j ' — ' o ' ² ' j ' — ' — ' μ ' » . ² ' μ w | ' , ' — ' — ' © @ — ' — ' â³ μ ' — ' w Q ² ' j ' — ' — ' » ' μ ' @ ' » ' » ³ ' o ' o ' μ — ' ² ' — ' — ' — ' p ' » ' ' » ' » ' © ' qº ' ² ' — ' — ' — ' ² ' — ' — ' « ' Q ' œ — ' — ' — ' @ ² , j — ' Q V ! ñ œ ' — ' — ' @ — ' — ' m ç Ÿ ' ' © ' » ' o ' o ' — ' — ' o a ' j ' « ' @ — ' — ' — ' μ ' | ' — ' — ' — ' — ' — ' œº » © , ' — ' — ' ± . © ' »º — ' — ' ± — ' μ Q « « ž μ μ ³ ' j ' μ — ' j ' — ' — ' » ' © ' » ' © ' | â « ' — ' — ' ² , j — ' Qº ± ' « 1 © — ' — ' É ' m μ™ m μ™ | ' m ç Ÿ ' ' © ' »º m μ™º q @ ³ ' TM — ' — ' — ' @ μº — ' — ' — ' — ' — ' w j ' — ' — ' œ » w Q W ' n — ' — ' — ' — ' — ' n — ' j ' « ' @ — ' — ' — ' yº — ' — ' ²º — ' ²º TX ' ² — ' — ' — ' — ' w Q ' m™º « « ž j μ — ž μ μ ³ ' — ' — ' — ' ³ ' j ' — ' — ' » ' © ' » ' © ' — ' — ' — ' , 1 ½º ' @ ² , j — ' Q

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

© 2014 Bayer CropScience



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Fenpyroximate 5% SC**(Acaricide)**

Fenpyroximate 5% SC is an acaricide used for the control of Yellow mites on Chillies, Red spider mite, Pink mite and Purple mite on Tea.

Chemical Composition:	Ingredient	%W/W
Anti-freezing agent (Propylene glycol)	Anti-freezing agent	5.0 % w/w
Defoamer (Dimethyl polysiloxane)	Defoamer	0.25 % w/w
Dispersing agent (Polyoxyethylene styryl phenyl ether and polyoxyethylene alkyl aryl ether)	Dispersing agent	6.0 % w/w
Distilled water	other ingredient	Q.S. to make 100% % w/w
Fenpyroximate a.i. content	active ingredient	5.0 % w/w
Preservative (1,2-benzisothiazin-3-one)	Preservative	0.1 % w/w
Thickener (Xanthum Gum) (Heteropolysaccharide)	Thickener	4.0 % w/w
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water Dilution in water (Litre)	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI a.i. (g)	Formulation Formulation (mL)					
Tea	Red spider mite Pink mite Purple mite	30-60	600-1200	400	Vegetative stage.	48 hours after spraying.	7	2-3 foliar sprays depending on pest pressure
Chilli	Yellow mite	15-30	300-600	500-750	Vegetative stage.	48 hours after spraying.	3	2-3 foliar sprays depending on pest pressure

Note:-**Good Practices relating to pesticide application techniques****Direction of Use**

Use recommended dose. Dilute the formulation in required quantity of water.

Weather Condition(in general):

Do not spray during rainy, windy and hot sunny period.

Soil & Water(where ever applicable):

This is foliar application product, so hardly have any relation to soil. Avoid using near aquaculture.

Crop Stage for application:

Vegetative stage.

Application Technique

Foliar spray using high volume sprayers.

Equipment Type:**Knapsack sprayer, foot sprayer, compression knapsack sprayer, compression knapsack battery sprayer or ASPEE - HTP sprayer.****NozzleType:****Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "**

=

Dosage:

15-30 g a.i./ha (300-600 mL formulation/ha) in 500-750 litres of water on Chilli and 30-60 g a.i./ha (600-1200 mL formulation/ha) in 400 litres of water on Tea.

Instruction for Mixing:

Re-entry Period

48 hours after spraying.

Manner and Time of Application

Application Timing

Should be applied preferably in the morning or evening hours.

Frequency

2-3 foliar sprays depending on pest pressure

Stages during which pesticides should not be applied on any crop :

=

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

Symptoms Of Poisoning

Symptoms may be combination of dizziness, headache, nervousness, anxiety, salivation tremors, convulsions, allergy, localized skin effect like itching, nausea, slight skin erythema and hyperactivity.

First Aid

1. If ingested : If person is conscious and alert, rinse mouth with water and give 1 or 2 glass of water to drink. Never give anything by mouth to an unconscious person. Call the doctor. Do not induce vomiting until and unless desired by attending physician. 2. If inhaled : Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration, preferably with mouth to mouth breathing. Consult a physician. 3. In case of skin contact : Wash thoroughly the affected part of body with soap and plenty of water including hairs and under fingernails. If pain, redness or irritation persists, consult a doctor. All the contaminated clothes should be segregated and thoroughly washed with soap and water separately before use. 4. In case of eye contact : Rinse gently with plenty of water for at least 15 minutes. If pain, irritation, redness or photophobia persists, consult eye specialist.

Antidote

There is no specific antidote. Hence, symptomatic and supportive therapy should be given.

Phytotoxicity

Fenpyroximate 5% SC is not phytotoxic when used as per recommendations.

Storage Conditions

1.it shall be the duty of manufacturers, formulators of insecticides and operators to dispose The packages containing the Acaricide shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles particularly food articles or shall be kept in separate almirahs under lock and key depending on the quantity and the nature of the Acaricide.

2. The store room should be well built, well lit, ventilated and sufficient in dimensions. The condition of the store should be dry and cool.

Safety Precaution:

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

1) Avoid inhalation and contact with skin and eyes. 2) Keep away from children, food stuff, animal feed, heat and flame, Store in cool and dry place under lock and key. 3) Do not mix the insecticides with bare hands. 4) Do not eat, drink, smoke or chew tobacco while spray. 5) Spray the insecticide only in the direction of wind blowing and not against it so as to prevent the setting of the drift on body. 6) The user should use full protective clothing with rubber gloves, boots. Face should be covered with face shield or dust mask and body should be covered with an overall or rubber apron and hood or hat.

Cautionary Statement:

1. not to be used for any purpose other than subscribed on label & leaflets.

2. destroy container after use as directed on leaflet.

3. do not use in those situations if there is possibility of harming bees & fish, livestock, game, wild birds & animals.

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufactures, formulators of insecticides and operators to dispose packages or surplus material and washings in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

M/s Rallis India Limited, C-5/C-6, MIDC Industrial Area,Phase-III,
Shivani, Akola . 444 104, MAHARASTHRA
Karnataka

M/s Rallis India Limited, Plot No. 2808, G.I.D.C. Estate,
Ankleshwar - 393002, Dist. Bharuch, GUJARAT
Karnataka

M/s Rallis India Limited, Plot No. 3301, GIDC Industrial Area, Dist.
Bharuch, Ankleshwar-393 002, GUJARAT
Karnataka

Rallis India Ltd Plot No. C44, Chemical Zone, Dahej Taluka-
Vagra, District- Bharuch - 392130, GUJARAT
Karnataka

Manufactured by

Rallis India Limited
Rallis India Limited, IIIrd Block, IIInd Floor, KSCMF Ltd., Building, 8,
Cunningham Road, Bangalore - 560052 Rallis India Limited
156/157, Nariman Bhavan, 15th Floor, 228, Nariman Point,
Mumbai - 400021 Bangalore Karnataka 560052

Customer Care Details:

Address: Rallis India Limited, IIIrd Block, IIInd Floor, KSCMF Ltd.,
Building, 8, Cunningham Road, Bangalore - 560052,
Email: kjsrivalli@rallis.co.in,
Contact No: 09810991089

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "