

Primary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Imazethapyr 70% WG

BRAND NAME: -

Chemical Composition :		Content % w/w
Total	-	100.00 % w/w
Imazethapyr (Active Ingredient)	Active Ingredient	70.00 % w/w
Sodium Alkyl naphthalenesulfonate blend	Other ingredient	2.00 % w/w
Alkylated naphthalene sulfonate formaldehyde polymer salt	Other ingredient	5.00 % w/w
Kaolin clay	Other ingredient	Q.S to make 100.00% % w/w
Total:		100.00%

Please refer leaflet for details

Antidote

there is no specific antidote. treat the patient symptomatically.

Manufactured by

???. ??, ?? ?? ???. ??, ?. ????? ??, ?????, ???????
 ?????-?????

Route No-2, Km 47.3, P.O. Box -243, Manati, Puerto Rico 00674

Supplied by

BASF Agricultural Products De Puerto Rico
 Route No-2, Km 47.3, P.O. Box -243, Manati, Puerto Rico 00674

Imported and Repacked by

Imazethapyr 70% WG

o² j^a ® ZS³ i

Broad Spectrum Systemic Selective Post Emergence Herbicide

»q®³ TM^o m© | » 'qñ®0-» -' -'® œ©³ TM[»] -' : μ'œ-ô«

° çœ' -' ®œ° ' «

Keep out of reach of children.

° ® - | ' ¥ð | ' i TM[»] ® ' n TM^o «

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Net Content / Volume _____

Max. Retail Price _____

Customercare _____

q®³ TM^o m© | » 'qñ®0-» -' -'® œ©³ TM[»] -' : μ'œ-ô«

o² j^a ® ZS³ i

-Á

° çœ' -' ®œ° ' «

-μ _____ 100.00 % w/w

o² j^a ® ZS³ i _____ 70.00 % w/w

S S -fl _____

° j³ 2³ _____ mð³ ' S S - _____ 2.00 % w/w

mý-om - ® μ - _____

° y⁻ -æ' 2³ _____

mý-μ æ' - ® μ - _____ mð³ ' S S - _____ 5.00 % w/w

° y⁻ -æ' 2³ μ | » Oⁱ _____

® μ 2³ ° yæ _____

- (μ - â μ ! _____ mð³ ' S S - _____ â³ ! w^o ! TSS³ ' ± ' ± ' _____ w/w

3 TM^o - TSS³ "

- p³ ' æ' 1³ - - - - - » » Q² ' Á - ' q®œ³ μ ¼³ - ° ' - ' Q

° ' w w^o ' w^o ' m^o - yœ μ^o | â æ^o b^o ' ð³ ' æ' - _____

æ' - ! UŽ - w^o ' WZ ! VŽ ® ! { ! ° â^o ' UWŽ² - © Ž ð³ ' æ' - ZSSYZW

®© ' - © | _____

° ' w w^o ' w^o ' m^o - yœ μ^o | â æ^o b^o ' ð³ ' æ' - _____

æ' - ! UŽ - w^o ' WZ ! VŽ ® ! { ! ° â^o ' UWŽ² - © Ž ð³ ' æ' - ZSSYZW

n³ © | ' ® - ® - TM^o - © | _____

Big Label

(> 250 mL Primary and Secondary Package Label)

Read leaflet before use:

Common Name:

Imazethapyr 70% WG

BRAND NAME: -

Crops and Pest Groups: -

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

Imazethapyr 70% wg is selective herbicide which provides control of grasses and broad leaf weeds in soybean.

Re-entry:

after 24 hours of spray.

Chemical Composition :

		Content % w/w
Total	-	100.00 % w/w
Imazethapyr (Active Ingredient)	Active Ingredient	70.00 % w/w
Sodium Alkyl naphthalenesulfonate blend	Other ingredient	2.00 % w/w
Alkylated naphthalene sulfonate formaldehyde polymer salt	Other ingredient	5.00 % w/w
Kaolin clay	Other ingredient	Q.S to make 100.00% % w/w

Total:100.00%

Safety Precautions:

Imazethapyr 70% WG

o² j^a @'ZS' 'j'øj³ j

Broad Spectrum Systemic Selective Post Emergence Herbicide

»q@3 TM' m© | » 'qñ@0- '» -' -'® œ©' 3 TM' » -' : μ' œ-0«



° çœ' -' ®œ°° '«

Keep out of reach of children.

° -' | '¥ð! 'j' TM' ®' n TM' ° '«

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Net Content / Volume _____

Max. Retail Price _____

q@3 TM' ° '®μ' ® -' ®E-

o² j^a @'ZS' 'j'øj³ j

'-A

'-A

o² j^a @'ZS' 'j'øj³ j' w-œ-« '®œ' -' -' »' j' ° -' q@3 TM' ° 3 ° -' 2 'š ° 'j' œE' ® -' μ' ®œ' -' -' -' 3 ' -' -' μw -' 3 ' j' ©' »Q

Y Ê- -' -' UWš °° «Q

' ° 3 -' -' ° œ- -

.....2 ' ± "±

-μ - 100.00 % w/w

o² j^a @'ZS' 'j'øj³ j' ° 3 'š ° - 70.00 % w/w

š ° -fl 2.00 % w/w

° j' 3 2 ' mō3 'š ° - 2.00 % w/w

mý-œ-® μ - ° ý- -œ' 2 ' 5.00 % w/w

m ý-μ š ° -® μ - mō3 'š ° - 5.00 % w/w

° ý- -œ' 2 ' μ j' » o j' ®μ 2 ' ° ý °

-{ μ - âμ! mō3 'š ° - â3 !w! 'TSS' ± '± 'w/w

3 TM' - TSSISS'

(1). BASF India Limited, 226/1, 226/2 GIDC Industrial Estate,
Panoli- 394 116, Gujarat
Gujarat

(2). Plot 5008, GIDC Estate, Ankleshwar, Dist. Bharuch, Gujarat .
393 001
Gujarat

(3). Unit No. 4 ,Village Nimbua, P. O Mubarakpur - 140201, Tehsil
Dera Bassi, Distt. Mohali , Punjab
Punjab

(4). SIDCO, Lane 2, Phase 1, Bari-Brahamana, Dist - Purmandal, J
& K 181133

(1).

(2).

(3).

(4). ?????, ??? 2, ??? 1, ???- ?????, ??, ?????, ?? ??? ??
181133

Customer Care Details

Address: BASF INDIA LIMITED, BASF INDIA LIMITED, Plot No. 37,
Chandivali Farm Road, Chandivali, Andheri (East) Mumbai .
400072,
Contact No: 022-67127827
Email: anirban.gangopadhyay@basf.com

» « ± μ · · ·

© - BASF INDIA LIMITED, BASF INDIA LIMITED, Plot No. 37,
Chandivali Farm Road, Chandivali, Andheri (East) Mumbai .
400072
μ - anirban.gangopadhyay@basf.com
Contact No: 022-67127827

Safety Pictograms:



Imazethapyr 70% WG

(Broad Spectrum Systemic Selective Post Emergence Herbicide)

Imazethapyr 70% WG is selective herbicide to be used in soybean for control of grasses and broad leaf weeds.

Chemical Composition:	Ingredient	%W/W
Alkylated naphthalene sulfonate formaldehyde polymer salt	Other ingredient	5.00 % w/w
Imazethapyr (Active Ingredient)	Active Ingredient	70.00 % w/w
Kaolin clay	Other ingredient	Q.S to make 100.00% % w/w
Sodium Alkyl naphthalenesulfonate blend	Other ingredient	2.00 % w/w
Total	-	100.00 % w/w
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water Liter (Water)	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI grams	Formulation grams					
Soybean	Cyperus rotundus, Echinochloa spp, Dinebra arabica, Digitaria spp, Brachiaria mutica Commelina benghalensis Commelina communis Euphorbia geniculata Cyanotis axillaris	70	100	500	to be applied when weeds are 2-3 leaf stage	After 24 hours of the spray	56 days	1 Spray

Note:-

Good Practices relating to pesticide application techniques

Direction of Use

Mix thoroughly required quantity of Imazethapyr 70% WG, Surfactant (Cyspread) and Ammonium Sulfate in a bucket containing water with adequate stirring with rod or stick. Add the remaining quantity of water as specified through agitation.

Weather Condition(in general):

Should be applied preferably in the morning and evening. Do not spray against wind direction

Soil & Water(where ever applicable):

To be properly tilled and Moisture level maintained before application.

Crop Stage for application:

to be applied when weeds are 2-3 leaf stage

Application Technique

Spray Imazethapyr 70% WG using high volume sprayer like Knapsack sprayer/ Foot sprayer fitted with flat fan nozzle

Equipment Type:

Knapsack sprayer/ Foot sprayer

NozzleType:

flat fan nozzle

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Dosage:

Soybean- 100 ml formulation per ha in 500 liters of water

Instruction for Mixing:

-

Re-entry Period

After 24 hours of the spray

Manner and Time of Application

Imazethapyr 70 % WG is an early post emergence herbicide to be applied when weeds are 2-3 leaf stage.

Application Timing

Imazethapyr 70 % WG is an early post emergence herbicide to be applied when weeds are 2-3 leaf stage.

Frequency

1 Spray

Stages during which pesticides should not be applied on any crop :

=

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

-Toxic to fish, hence should not be used near aquaculture.

Symptoms Of Poisoning

Irritation to skin and eyes may occur. Headache, nausea, vomiting etc may occur.

First Aid

In case of skin contact remove contaminated clothes immediately. Wash the contaminated skin thoroughly with soap and plenty of water. If inhaled: Keep patient calm, remove to fresh air. In case of eye contact: wash affected eyes for at least 15 minutes under running water with eye lids held open. If ingestion: Rinse mouth immediately and then drink plenty of water. Seek medical attention.

Antidote

There is no specific antidote. Treat the patient symptomatically.

Phytotoxicity

Not phytotoxic when used at recommended dose

Storage Conditions

1. The packages containing the herbicides should be stored in original containers in separate rooms or premises away from the rooms or premises used for storing other articles particularly articles of food or shall be kept in separate almiraha under lock and key depending upon the quantity and nature of herbicide.
2. The rooms or premises meant for storing the herbicide shall be well built, airy, well lit and ventilated and of sufficient dimension to avoid contamination with vapour.

Safety Precaution:

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. Do not mix with bare hands.
2. The user should use full protective clothing, which include rubber gloves, boot, face shield and overall or rubber apron or hat.
3. Keep away from foodstuffs, empty foodstuffs containers and animals food.
4. Do not smoke, drink, eat and chew anything while spraying. Wash the affected areas before eating, drinking or smoking. Avoid contamination of environment and water.

Cautionary Statement:

not to be used other than the specified crop.

Toxic to fish, hence should not be used near aquaculture.

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufactures, formulators of insecticides and operators to dispose packages or surplus material and washings in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

BASF India Limited, 226/1, 226/2 GIDC Industrial Estate, Panoli-394 116, Gujarat Maharashtra
Plot 5008, GIDC Estate, Ankleshwar, Dist. Bharuch, Gujarat . 393 001 Maharashtra
SIDCO, Lane 2, Phase 1, Bari-Brahamana, Dist - Purmandal, J & K 181133 Maharashtra
Unit No. 4 ,Village Nimbua, P. O Mubarakpur - 140201, Tehsil Dera Bassi, Distt. Mohali , Punjab Maharashtra

Imported and Repacked by

BASF INDIA LIMITED
BASF INDIA LIMITED, Plot No. 37, Chandivali Farm Road, Chandivali, Andheri (East) Mumbai - 400072 MUMBAI Maharashtra 400072

Customer Care Details:

Address: BASF INDIA LIMITED, BASF INDIA LIMITED, Plot No. 37, Chandivali Farm Road, Chandivali, Andheri (East) Mumbai . 400072,
Contact No: 022-67127827
Email: anirban.gangopadhyay@basf.com

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "