

Primary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

N.P.V. of Helicoverpa armigera 2.0% AS

BRAND NAME: -

Chemical Composition :

| Content % w/w | |
|--|---|
| Water | Water Q.S. to make - |
| Total | Total 100.00% W/W |
| NPV of Helicoverpa Armigera in Hosttissue Residues | Armigera in Hosttissue Residues 2.00% AS |
| POBs count/ml | POBs count/ml 1x10 ⁹ per ml.min. |
| Glycerine | Glycerine 5.00% - |
| Total: 100.00% | |

Please refer leaflet for details

Antidote

treat symptomatically.

Manufactured by

R B Herbal Agro
M/s R.B. HERBAL AGRO Herbal House, 120, Savarkar Nagar,
Satana 423301, Distt:- Nashik (M.S.) Nashik Maharashtra

Premises Address

(1). Gat No.143, Chaugaon, Tal- Satana, Distt:-
Nashik-423301(Maharashtra)
Maharashtra

Customer Care Details

N.P.V. of Helicoverpa armigera 2.0% AS

» μ— . @! n 2 T M r ° — ð³ â μ ³ ° μ » | ° ° — π ° ° z U ° W °
W ° ! f w - ! @ ! ° ! fl

Contact Insecticide

° 3 ° — π — ° —



° ° ° | ° ° ¥ 0 | ° ° T M » ° ° n T M ° ° ° ° ° ° ° °

Keep in cool, dry place away from heat and open flame.

| | | |
|----------------------|----------------------------|-------|
| Regn. No. | ° j ! ° ! | _____ |
| Mfg. Lic. No. | 2 _ 3 ! ° μ ! ° ! | _____ |
| Batch No. | ° œ° ! | _____ |
| Mfg Date | 2 _ 3 ! ° @ ! | _____ |
| Exp. Date | ¿ ñ² ° » — ° ° ° ° ° ° ° ° | _____ |
| Net Content / Volume | ° ± ñ ³ ° _ | _____ |
| Max. Retail Price | m n — ° 2 ° ° ° ° ° ° 2 y³ | _____ |
| Customercare | | _____ |

q ð³ T M r ° ° ° ° μ ° — ° ° E° -

» μ— . @! n 2 T M r ° — ð³ â μ ³ ° μ » | ° ° — π ° ° z U ° W °
W ° ! f w - ! @ ! ° ! fl

° 3 ° — π — ° —

| | | |
|----------------------|----------------------------|-------|
| Regn. No. | ° j ! ° ! | _____ |
| Mfg. Lic. No. | 2 _ 3 ! ° μ ! ° ! | _____ |
| Batch No. | ° œ° ! | _____ |
| Mfg Date | 2 _ 3 ! ° @ ! | _____ |
| Exp. Date | ¿ ñ² ° » — ° ° ° ° ° ° ° ° | _____ |
| Net Content / Volume | ° ± ñ ³ ° _ | _____ |
| Max. Retail Price | m n — ° 2 ° ° ° ° ° ° 2 y³ | _____ |
| Customercare | | _____ |

XISS ° ° ° ° z

3 T M r ° — TSSISS °

μ ¼ ° ° ° ° μ j ° — ° ° ° °

q ñ ° ° —

° ° ° ° ° ° ° ° ° °

(1).

» — ° ° ± μ ° ° ° ° ° °

R.B. HERBAL AGRO
Herbal House, 120, Savarkar Nagar, Satana 423301, Distt: -
Nashik (M.S.)

n 1° !' »° μ' w
»° μ' » q° Ž 120Ž° . '—'—™Ž° ©— ' 423301Ž i μ -'ž' - ° —
fW? !w? !fl

Safety Pictograms:



Secondary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

N.P.V. of *Helicoverpa armigera* 2.0% AS

BRAND NAME: -

Crops and Pest Groups: Chickpea- Chick pod borer (*Helicoverpa armigera*)

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For Agriculture use.

Re-entry:

12 Hrs.

Chemical Composition :

Content % w/w

| | | |
|---|--|-------------------------------|
| Water | Water | Q.S. to make - |
| Total | Total | 100.00% W/W |
| NPV of <i>Helicoverpa Armigera</i> in Hosttissue Residues | <i>Armigera</i> in Hosttissue Residues | 2.00% AS |
| POBs count/ml | POBs count/ml | 1x10 ⁹ per ml.min. |
| Glycerine | Glycerine | 5.00% - |

Total: 100.00%

Safety Precautions, Symptom of Poisoning and First Aid :Please refer leaflet for details.

Antidote Statement:

treat symptomatically.

Cautionary statement:

N.P.V. of *Helicoverpa armigera* 2.0% AS

» μ— . @I n 2 T M r ' — ' 0 3 â μ 3 ' ' @ μ » | ' ° ° ' — α ' ' ž U ' ' w ' w ' ! f w - ! @ ! : ! f l

Contact Insecticide

@ , 3 ' — α — , —



° ° ' | ' ' ž U ' ' w ' ' — ' 0 3 â μ 3 ' ' @ μ » | ' ° ° ' — α ' ' ž U ' ' w ' ' ! f w - ! @ ! : ! f l

Keep in cool, dry place away from heat and open flame.

| | |
|----------------------|-------|
| Regn. No. | _____ |
| Mfg. Lic. No. | _____ |
| Batch No. | _____ |
| Mfg Date | _____ |
| Exp. Date | _____ |
| Net Content / Volume | _____ |

q®³ TM' o ' ®»μ' ® — ®E-

» μ— . @I n 2 T M r ' — ' 0 3 â μ 3 ' ' @ μ » | ' ° ° ' — α ' ' ž U ' ' w ' w ' ! f w - ! @ ! : ! f l

— 1 ' q®³ TM' — ' μ w

» μ— . @I n 2 T M r ' — ' 0 3 â μ 3 ' ' @ μ » | ' ° ° ' — α ' ' ž U ' ' w ' w ' ! f w - ! @ ! : ! f l

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For Agriculture use.

Re-entry:

12 Hrs.

Chemical Composition :

| | | |
|---|--|-------------------------------|
| Water | Water | Q.S. to make - |
| Total | Total | 100.00% W/W |
| NPV of <i>Helicoverpa Armigera</i> in Hosttissue Residues | <i>Armigera</i> in Hosttissue Residues | 2.00% AS |
| POBs count/ml | POBs count/ml | 1x10 ⁹ per ml.min. |
| Glycerine | Glycerine | 5.00% - |

Total: 100.00%

Safety Precautions, Symptom of Poisoning and First Aid :Please refer leaflet for details.

Antidote Statement:

treat symptomatically.

Cautionary statement:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Max. Retail Price _____

Manufactured by

R B Herbal Agro
M/s R.B. HERBAL AGRO Herbal House, 120, Savarkar Nagar,
Satana 423301, Distt: - Nashik (M.S.) Nashik Nashik Maharashtra
Premises Address

(1). Gat No.143, Chaugaon, Tal- Satana, Distt:-
Nashik-423301(Maharashtra)
Maharashtra

Customer Care Details

R.B. HERBAL AGRO
Herbal House, 120, Savarkar Nagar, Satana 423301, Distt: -
Nashik (M.S.)

Safety Pictograms:



(1).

(1). Gat No.143, Chaugaon, Tal- Satana, Distt:-
Nashik-423301(Maharashtra)
Maharashtra

Customer Care Details

R.B. HERBAL AGRO
Herbal House, 120, Savarkar Nagar, Satana 423301, Distt: -
Nashik (M.S.)

Safety Pictograms:



(1).

»—'«~ ± μ'

n '1° !' » μ' w

»° μ' » q° Ž 120Ž°™Ž° ©— : 423301Ž j μ - ž - ° —

fW? !W? !fl

» μ—, @In 2™ —ô³âμ³@μ»|°°—«'ž'U' 'w'w'!fw-!@!- !f|

fl @ 3'—«—, —fl

1 10 ml.min

» μ—, @In 2™ —ô³âμ³@μ»|°°—«'ž'U' 'w'w'!fw-!@!- !f| —«—, —»|°²» μ—, @In 2™ —ô³âμ³@μ»|°°—«'ž'U' 'w'w'!fw-!@!- !f| —«—, —»|C
 3»'œ'—'°μ'—'—° —'—'': μ'—«'—'—«'—'©'»'Q

° 3 —'°œ'—

| | | |
|---|------------------------|--------------------|
| âμ°'— | âμ°'— | XISS'—ž |
| w-!@!y»'y Ó'» μ—, @In 2™ —'o—'» «p°' w-!@!y»'y Ó'» μ—, @In 2™ —'o—'» «p°' USS'—AS | | |
| ° 3 | ° 3 | |
| @!{ !° !w'!°° â³'—'²!μ | @!{ !° !w'!°° â³'—'²!μ | 1x10^9 per ml.min. |
| —μ | —μ | TSSISS'—W/W |
| @'' | @'' | @³ b@²'—ž |
| | | 3™ — TSSISS'—±'±' |

| ° μ' | —«'—'—² | °â³'—² | | @—'—² μ«' | ° μ'm' a | @—'—μ | @ l'° μ | n . |
|------|---------------------------|------------------------------------|----------------|--------------|----------|-------|---------|----------|
| | | ° !'©ñ w-!@!- !» μ— . @In 2™ | ° 'œ'— ²! μ | | | | | |
| œ- | œ- 'ÿ«— f» μ—'—'w²™ fl | » μ—, @In 2™ —'U' 'w'w'! | UXS' w' !vμ | YSS μ«' | - | - | - | j° 'Á' © |

Note:-

g@³™ — μw —«'—

œ'—'YSS' μ«'@—'²' UXS' ²!μ !w-!@!- !'U' 'w'w'!š μ—'TX' «'—'m@' μ'@'«'°°' ©-³'—³²'—°²³'°š μ'—'ÿ|—'—
—Q

2°²'—' a ©

° —'n°² —

2 «'|'—'©—

—°'±'—'—'²«

° μ'm' a

n . «'—'©—

-

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

_____:

1!—x,—@m—@°23Z~μ~3~Y~x~—~j~»~°~°~°~|~ñ~æ~—~°~@~°~°~αQ2!~!~TM~»~a~°~°~—~x,—~—~2~μ~wQ3!~ñ~3~—
q® TM° «Z~2~—~|~ñ~Y~Ë—~q@—~°~°~mõ3~—~x,—~—~Y~Ë—~—~μwq® TM~—~—~°~°~®~μ~°~»~°~°~@~—~|~ñ~j~x~3~°~°~—~—~
š~μ~°~°~mçY~°~°~»~°~°~wQ4!~°~æ~°~3~—~x,—~—~2~—~—~—~a~°~°~3~°~°~»~°~°~@~—~2~Z~°~μ~—~±~2~Z~j~»~j~μ~°~°~—~°~°~«~1~@~—~—~°~°~±~—~—~
»~Z~·~·~»~—~Y~|Q5!~q® TM~@~|~—~°~°~1°~°~¼~n~°~—~@Ë~—~q® TM~—~—~°~°~æ~°~w~j~°~2~°~°~°~°~«~°~°~°~æZ~3~°~°~š~μ~°~3~—~μ~2~°~°~—~|~ñ~
°~2~°~°~°~w~Z~»~j~°~°~3~°~°~x~®~°~°~2~μ~°~»Q6!~°~æ~μ~—~|~ñ~3~TM~—~«~°~°~°~°~@~Z~°~°~3~°~°~—~—~—~Q

_____:

Tflμ°μ°|ñ°æ°®°j°m±°»Z°q°°°±õ°mõ3°3j°~2°q®TM~μ°3°~»!°fufi°æ°°°~«°°°q®TM~°°æ°m~—~—~x~
—~—~

_____:

1~3~»~—~x,—~—~2~|@{Zy~°~°~x~°~|~ñ~2~μ~x~—~—~y~3~°~»~TM~—~@~3~j~°~°~3~°~3~j~μ~°~°~«~1~°~°~—~—~—~—~μww~°~°~¼°~°~°~—~°~°~
®~j~°~3~°~m~n~j~°~°~2~°~°~|~ñ~n~—~—~—~@~x~—~—~°~°~Q
2!°q®TM~—~3~°~»w°®~j~°~°~—~—~μ~2~°~°~—~Y~Ë~°~—~q~—~°~°~®~3~TM~—~—~3~j~°~°~—~°~°~Q
3!°®~j~°~°~°~@~Ë~°~°~°~n~°~«~°~°~°~°~j~2~°~°~—~—~—~æ~TMË°~°~«~Q

qñ®«~°~°~°~

qñ®«~—

»~°~«~°~±~μ~°~°~°~

n°!°!»°°μ°w
»°μ°°q°Z120Z°°°—~—~—~TM~Z°°@~°°423301Z~j~μ~°~Z~°~°~—~°~°~fw°!w°!fl

°~¼°~æ°°μ~®~



Caution: Í Not to Be Used on Crops Other Than Specified on This Label/Leaflet! "

N.P.V. of Helicoverpa armigera 2.0% AS

(Contact Insecticide)

1 10 ml.min

Nuclear Polyhedrosis Virus of Helicoverpa armigera-2% AS is a biological insecticides contains spores of Nuclear Polyhedrosis Virus of Helicoverpa armigera-2% AS, a naturally occurring entomopathogenic virus that infects the insect pests of chick pea

| Chemical Composition: | Ingredient | %W/W |
|--|---------------------------------|-------------------------------|
| Glycerine | Glycerine | 5.00% - |
| NPV of Helicoverpa Armigera in Hosttissue Residues | Armigera in Hosttissue Residues | 2.00% AS |
| POBs count/ml | POBs count/ml | 1x10 ⁹ per ml.min. |
| Total | Total | 100.00% W/W |
| Water | Water | Q.S. to make - |
| Total: | | 100.00% w/w |

Recommendation

| Crop(s) | Common Name of Pest | Dosage/HA | | Dilution in Water Litre. | Crop Stage | Re-entry Period | Pre Harvest | Frequency |
|----------------|--|--|-----------------------|---------------------------------|-------------------|------------------------|--------------------|------------------|
| | | AI N.P.V. of Helicoverpa armigera | Formulation ML | | | | | |
| Chickpea | Chick pod borer (Helicoverpa armigera) | Helicoverpaarmigera-2%AS | 250ML | 600Litre | - | - | - | as required |

Note:-

Good Practices relating to pesticide application techniques

Direction of Use

Dissolve 250 mls of NPV-2% AS in 600 ltrs of water and spray the solution twice at an interval of 15 days preferably in the morning and evening hours.

Weather Condition(in general):

Clear Sky

Soil & Water(where ever applicable):

Any Type

Crop Stage for application:

-

Application Technique

-

Equipment Type:

Rocking or knapsack sprayer, hand compressor sprayer, foot sprayer, motorized knapsack sprayer

NozzleType:

Hollow cone nozzle

Dosage:

Caution:Í Not to Be Used on Crops Other Than Specified on This Label/Leafletí "

250 mls of NPV-2% AS in 600 ltrs of water

Instruction for Mixing:

-

Re-entry Period

-

Manner and Time of Application

preferably in the morning and evening hours.

Application Timing

spray the solution twice at an interval of 15 days

Frequency

as required

Stages during which pesticides should not be applied on any crop :

=

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

-

Symptoms Of Poisoning

Headache, Nausea, Vomiting, irritation & allergy may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to 15 minutes. 4. Remove the patient into fresh air.

Antidote

Treat symptomatically.

Phytotoxicity

No phytotoxic at Recommended Dose level.

Storage Conditions

1. The Packages containing the Bio control-insecticide shall be stored in separate room or premises away from the rooms or premises used for storing other articles particularly articles of food or shall be kept in separate almirahs under lock and key depending upon the quantity and nature of the Bio control-insecticide.
2. The rooms or premises meant for storing the Bio control-insecticide shall be well built, dry, well lit and ventilated and of sufficient dimension to avoid contamination with vapour.

Safety Precaution:

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothings which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

Cautionary Statement:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Disposal of used containers, washing from equipment and surplus material :

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

- (1) It shall be the duty of manufactures, formulators of insecticides and operators to dispose packages or surplus material and washings in a safe manner so as to prevent environmental or water pollution.
(2) The used packages shall not be left outside to prevent their re-use
(3) The packages shall be broken and buried away from habitation.

Manufacturer Premises :

Gat No.143, Chaugaon, Tal- Satana, Distt:-
Nashik-423301(Maharashtra)
Maharashtra

Manufactured by

R B Herbal Agro
M/s R.B. HERBAL AGRO Herbal House, 120, Savarkar Nagar,
Satana 423301, Distt: - Nashik (M.S.) Nashik Maharashtra

Customer Care Details:

R.B. HERBAL AGRO
Herbal House, 120, Savarkar Nagar, Satana 423301, Distt: - Nashik (M.S.)

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "