

Primary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Sulfosulfuron 75% WG

BRAND NAME: -

Chemical Composition :

Content % w/w

Ethelene oxide . Á	Suspending aid	1.70 w/w
Propylene oxide		
Ammonium sulfate	Filler	q.s. w/w
Sulfosulfuron a.i.	Active ingredient	75.00 w/w
Sodium Naphthalene Sulfonate	Dispersant	2.50 w/w
Lignosulfonate	Dispersant	0.10 w/w
Tallow soap	Defoamer	0.20 w/w
Sodium dialkyl naphthalene Sulfonate	Wetting agent	0.10 w/w
3EO alkyl (C12-C15) ether sulfate	Wetting agent	4.20 w/w
Alpha . Álky (C10-C16)-omega-hydroxypoly (oxyethylene) mixture	Antifoam agent	0.10 w/w
Silicone defoamer	Defoamer	0.10 w/w

Total: 100.00%

Sulfosulfuron 75% WG

° y- ° y- - ZX- ' | øy³ ' i

Selective Herbicide

œ³ - ñ² - ' ®©. ' - , -



° çœ³ - ' ®œ³ ° ' « ' - -

Keep out of reach of children.

° - - | ' ' ¥ð' ' i TM» ' ®' n TM ° ' « ' - -

Keep in cool, dry place away from heat and open flame.

Regn. No. ' i ! - !

Mfg. Lic. No. 2 - 3 ! ' µ ! - !

Batch No. ° œ³ - !

Mfg Date 2 - 3 ! ' © !

Exp. Date ç ñ² ' » - ' - ' © ' ' -

Net Content / Volume , ' ± ' ž ñ³ © -

Max. Retail Price m - - ©² ' ' « ' ' ² y³

Customercare

q®³ TM ° ' ®µ ' ® - ' ®£-

- - - - -

° y- ° y- - ZX- ' | øy³ ' i

- - - - -

' ° 3 - - - ' œ³ - -

wª µ - ' ° ° ® ! TM wµ

n á ° o ! ž ® µ - ' ° °

y á ° o !

m² - 3 2 ' ° y- ' - µ ' ° °

° y- ° y- - ' ° 3 ' ° 3 ' © ñ .

© ñ

° ' | 3 2 ' - - a µ - ' ' ° ® ° µ

° y- ' - µ

µ ä ' - ° y- ' - µ ' ' ° ® ° µ

µ µ ° ®

° ' | 3 2 ' | pmy ' - - om ' - µ TM wµ µ

- - a µ - ' ° y- ' - µ

V ' p { ' my ' - - pµ ' ff TU . µ TM wµ µ

ž° TXfl oª ' ' ° y- ' - µ

wy ' - žmy ' - - om ' ff TS wµ - 2 ' ' wµ µ

ž° TYfk { 2 TM ' ' -

» á ° ® µ ' ° °

fy á ° w µ - fl

² á ° œ³

° µ - - ' | - 2 ' ' | - 2 ' ' ° °

3 TM - TSS!SS

Please refer leaflet for details

Antidote

No specific antidote, symptomatic and supportive therapy

Big Label

(> 250 mL Primary and Secondary Package Label)

Read leaflet before use:

Common Name:

Sulfosulfuron 75% WG

BRAND NAME: -

Crops and Pest Groups: -

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

for agricultural use

Re-entry:

24 hours after spraying

Chemical Composition :

Content % w/w

Ethelene oxide . Á	Suspending aid	1.70 w/w
Propylene oxide		
Ammonium sulfate	Filler	q.s. w/w
Sulfosulfuron a.i.	Active ingredient	75.00 w/w
Sodium Naphthalene Sulfonate	Dispersant	2.50 w/w
Lignosulfonate	Dispersant	0.10 w/w
Tallow soap	Defoamer	0.20 w/w
Sodium dialkyl naphthalene Sulfonate	Wetting agent	0.10 w/w
3EO alkyl (C12-C15) ether sulfate	Wetting agent	4.20 w/w
Alpha . Álkyl (C10-C16)-omega-hydroxypoly (oxyethylene) mixture	Antifoam agent	0.10 w/w
Silicone defoamer	Defoamer	0.10 w/w

Sulfosulfuron 75% WG

° y° ° y° - ZX° | øy³ |

Selective Herbicide

œ³ - ñ² - - ©. - , -



° çœ³ - - ©œ³ ° ' «

Keep out of reach of children.

° - - | - - ¥ð! | | TM» | ©ñ TM ° ' «

Keep in cool, dry place away from heat and open flame.

Regn. No. | | -!

Mfg. Lic. No. 2 - 3 | ! -!

Batch No. ° œ³ -!

Mfg Date 2 - 3 | ! |

Exp. Date 2 ñ² ' » - - - © ' -

Net Content / Volume , ± ñ³ © -

Max. Retail Price m - - ©² ' » « ' 2 y³

q®³ TM ° ' ®µ ' ® - - ®É-

° y° ° y° - ZX° | øy³ |

- - Á

- - Á

- 1 ' q®³ TM° - - µwQ

ÿÉ - - - % ' šœ³ ° «

° ° 3 - - - ° œ³ - -

..... 2 - - ± "±

wª µ - ° ° ® | TM° W | TIZS° w/w

n á° ° o!ž ®µ -

y á° ° o!

m² - 3 2 ° y° - - µ ' ®³ b©² ' w/w

° y° ° y° - - ° 3° ° 3° ©ñ° ZXISS° w/w

©ñ

° | 3 2 - - ° a µ - | ° ®° ° SXUS° w/w

° y° - -

µä - ° y° - - | ° ®° ° SITS° w/w

µµ ° ® | - 2 ' SIUS° w/w

° | 3 2 | | pmy° - oµ ' . œTM° W | SITS° w/w

- - ° a µ - ° y° - -

V° p° { ' my° - pµ° ffTU . œTM° W | WUS° w/w

ž° TXfl oª ' ° y° - -

° n p⁻ VSZVX' «-° «

n · «-° —

÷ μ α: y³ 2³ ÷ μ | j α: j μ⁻ a μTM qçœ² : μ⁻ 3 | 'vw' @ppžwœ @ @. 3 '—q@³ TM—' @»w» ° p o | m- 3 TMQ

÷ μ α⁻ -"÷ μ | Å α: j μ⁻ a — @— 3 ° Q

÷ μ α⁻ -"÷ μ | Å α: j μ

33.3g. formulation /ha. (25ga.i./ha.) in 200-250 Olit. of water

1° y⁻ ° y⁻ '— ZX' | øy³ j @—α:—μ² : fTVIX' !flY' μ α'@— 2 | μ—'mçY' ©'»' 2 μ o³ Q «2—æn 3 —° '— â pff !'

2 â° fl—μ² : fXSS' 2 | μ!fb° @— 2 | μ—'mçY' ©'»' 2 μ o³ Q° 'y³ : TSS' μ α'@— 2 '2 μ —'w—w—Ê² 'Y Ê—Q

UWS α

° y⁻ ° y⁻ '— ZX' | øy³ j @—α:—μ² : fTVIX' !flY' μ α'@— 2 | μ—'mçY' ©'»' 2 μ o³ Q «2—æn 3 —° '— â pff !'

2 â° fl—μ² : fXSS' 2 | μ!fb° @— 2 | μ—'mçY' ©'»' 2 μ o³ Q° 'y³ : TSS' μ α'@— 2 '2 μ —'TM»—°. p⁻ 30-35' «— :'

° «w—w—Ê² 'Y Ê—Q

n · «-° — ° 2 3

m → 2 — @° ° » | : 2 — 2 3 Q

w—m— 3 TM

'—' @ @. '—qTM— @ μTM— » » © »

. 1' — μ ¼'

° y⁻ ° y⁻ '— ° n : œ² fl— μ— » ° — © » Qm → μ 'n — ž © μTM— žn ' ° @— n — ž ° μ— 2 @. — õñ³ «Q

@ «®' n . 1 μ ©

° ç . — 2 — m— ° ' 3 TM— '— @' 3 » qñ® «®— . , â © — » » Q

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

qñ@«-’@’°’-

3-11ž ĩn p!; !° !ž @žm ĩ ©ž 396195

3 !@!vμ! μ 2 μ!

3 – pμ! Ó Ó’° μ 2 μ! 3 – Ó ° » r ° ž š @— 11’ ĩ ž

fĭ μř ú° pž 400 0522 ú° p’ 400052

ođ; 3 μ’ a ° oα’ž ú° ž ú² žĭ mđ; —fl 184121

đμ μ—! 3101/02ž ĩn p!; !° !žnđμ · žm ĩ ©’ >393002

2 !—! 225žm. ™@®’ ž° !» μ μ žĭ !@αı» μ žm ĩ ©

»—« ± μ’ · · ‘—

° ¼’ œ’ μ @-



Caution: ĩ Not to Be Used on Crops Other Than Specified on This Label/Leaflet ĩ”

Sulfosulfuron 75% WG

(Selective Herbicide)

Sulfosulfuron 75 WG is a selective early post-emergence herbicide used for the effective control of Phalaris minor in wheat crop. It is also control broad leaf weeds like Chenopodium album, Melilotus alba.

Chemical Composition:	Ingredient	%W/W
3EO alkyl (C12-C15) ether sulfate	Wetting agent	4.20 w/w
Alpha . Alkyl (C10-C16)-omega-hydroxypoly (oxyethylene) mixture	Antifoam agent	0.10 w/w
Ammonium sulfate	Filler	q.s. w/w
Ethelene oxide . Propylene oxide	Suspending aid	1.70 w/w
Lignosulfonate	Dispersant	0.10 w/w
Silicone defoamer	Defoamer	0.10 w/w
Sodium dialkyl naphthalene Sulfonate	Wetting agent	0.10 w/w
Sodium Naphthalene Sulfonate	Dispersant	2.50 w/w
Sulfosulfuron a.i.	Active ingredient	75.00 w/w
Tallow soap	Defoamer	0.20 w/w
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water Litre/ha.	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI g. a.i./ha.	Formulation gram/ha.					
Wheat	Phalaris minor, Chenopodium album, Melilotus alba	25.0	33.3	200-250	30-35 days after sowing	24 hrs	110 days	One time application

Note:-

Good Practices relating to pesticide application techniques

Direction of Use

1. Dissolve the entire contents of Sulfosulfuron 75 WG sachet (13.5 g) in six litres of water and mix thoroughly.
2. Then add the entire contents of cationics surfactant (500 ml) to the above solution and mix thoroughly with a stick.
3. Dilute the solution with 100 liters of water to uniformly spray one acre.

Weather Condition(in general):

Do not spray during rainy, windy and hot sunny period

Soil & Water(where ever applicable):

This Herbicide is for foliar application and has no adverse effect when used at recommended dose . Avoid using near aquaculture

Crop Stage for application:

30-35 days after sowing

Application Technique

Herbicide application using high volume sprayers and ASPEE-HTP power sprayer fitted with flat fan or flood jet nozzle.

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

Equipment Type:
~~Knapsack sprayer~~ with flat fan/ flood jet nozzle.

NozzleType:

Flat fan or flood jet nozzle

Dosage:

33.3g. formulation /ha. (25ga.i./ha.) in 200-250 0lit. of water

Instruction for Mixing:

1. Dissolve the entire contents of Sulfosulfuron 75 WG sachet (13.5 g) in six litres of water and mix thoroughly.
2. Then add the entire contents of cationic surfactant (500 ml) to the above solution and mix thoroughly with a stick.
3. Dilute the solution with 100 liters of water to uniformly spray one acre.

Re-entry Period

24 hrs

Manner and Time of Application

1. Dissolve the entire contents of Sulfosulfuron 75 WG sachet (13.5 g) in six litres of water and mix thoroughly.
2. Then add the entire contents of cationic surfactant (500 ml) to the above solution and mix thoroughly with a stick.
3. Dilute the solution with 100 liters of water to uniformly spray one acre at 30-35 days after sowing of Wheat.

Application Timing

Preferably in the morning and evening hours.

Frequency

One time application

Stages during which pesticides should not be applied on any crop :

Pre-emergence of weed

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

.

Symptoms Of Poisoning

The product may cause irritation to the eyes and skin, excessive salivation, diarrhoea, lacrimation, laboured breathing etc.

First Aid

If ingested: if person is conscious and alert, rinse mouth with water and give 1 or 2 glass of water to drink. Never give anything by mouth to an unconscious person. Call the doctor. Do not induced vomiting until and unless desired by attending physician.

If inhaled: Take the patient to fresh air and take care of body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration, Consult a physician.

In case of skin contact: Wash thoroughly the affected part of body with soap and plenty of water including hairs and under fingernails. If pain, redness or irritation persists, consult a doctor. All contaminated clothes should be segregated and thoroughly washed with soap and water separately before use.

In case of eye contact: Rinse gently with plenty of water at least for 15 minutes. If pain, irritation, redness or photophobia persists. Consult eye specialist

Antidote

No specific antidote, symptomatic and supportive therapy.

Phytotoxicity

The product is not phytotoxic when used as per recommendations.

Storage Conditions

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

1. The packages containing the herbicide shall be stored in separate rooms or premises, away from the rooms or premises used for storing other articles or shall be kept in separate almira's under lock and key depending upon the quantity and nature of the herbicide.
2. The rooms or premises meant for storing the herbicide shall be well-built, dry, well-lit and ventilated and of sufficient dimension.

Safety Precaution:

1. Sulfosulfuron should only be applied as liquid spray. Do not broadcast Sulfosulfuron. 2. Do not use a Mist blower (Power spray).
3. The spray man should use appropriate protective clothing like hat, long sleeved shirt, long pants and shoes.
4. Avoid contact with eyes and skin when handling the concentrates spray solution. Avoid inhalation of spray mist. Wash thoroughly with soap and water after handling. Do not eat, drink or smoke during application. Remove contaminated clothes and wash before reuse. Do not re-use the empty containers.
5. Wash hands & exposed skin with soap and water before meals and after work.
6. Dicot crops like mustard, gram etc. should not be grown at inter crop in wheat.
7. After spraying this herbicide, the sprayer and utensils should be thoroughly cleaned with plenty of soap and water.

Cautionary Statement:

- 1. It is toxic to honey bees. spray application should be avoided during the foraging period of bees.**
- 2. toxic to fish, aquatic organisms hence should not be used near water bodies, aquaculture or pisciculture.**
- 3. Not to be used for any purpose other than those subscribed on label and leaflets.**
- 4. Destroy the container after use as directed on leaflet.**
- 5. Toxic to earthworms and other beneficial arthropods.**

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufacturers, formulators of insecticides and operators to dispose packages or surplus materials and washing in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

3 - 11, G.I.D.C., Vapi, Gujarat-396 195
Maharashtra

Industrial Growth Centre, Samba, Jammu, (J & K) 184121
Maharashtra

Plot No. 3101/02, G.I.D.C, Ankleshwar, Gujarat-393 002
Maharashtra

S. no. 225, Village Gopipura, Ta. Halol, Dis. Panchmahal
Maharashtra

Manufactured by

UPL Limited
Uniphos House, Madhu Park, 11th Road, Khar (W), Mumbai 400
052 MUMBAI Maharashtra 400052

Customer Care Details:

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "