

Primary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Tebuconazole 6.7% Captan 26.9% w/w SC

BRAND NAME: -

Chemical Composition :

Content % w/w

Captan a.i.	active ingredient	26.9 % w/w
Tebuconazole a.i	active ingredient	6.7 % w/w
Polyarylphenyl ether phosphate	other ingredient	2.1 % w/w
Propylene glycol	other ingredient	5.9 % w/w
Xanthan Gum	other ingredient	0.25 % w/w
Water	other ingredient	q.s % w/w
Modified-dimethyl polysiloxane	other ingredient	0.6 % w/w
Sodium Naphthalene formaldehyde condensate	other ingredient	1 % w/w

Total: 100.00%

Please refer leaflet for details

Antidote

no specific antidote, treat symptomatically.

Manufactured by

ADAMA India Pvt. Ltd.
Plot No: DS-13, IKP Knowledge Park, Sy. No. 542/2, Genome Valley, Turkapally, Shameerpet, Ranga Reddy District Hyderabad Hyderabad Telangana 500078

Premises Address

Tebuconazole 6.7% Captan 26.9% w/w SC

Fungicide



Keep out of reach of children.

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Net Content / Volume _____

Max. Retail Price _____

Customercare _____

(1). S.No. 180/6, & 180/7, Khazipally (V), Jinnaram (M), Dist-
Medak, Telangana-502 319

(1).

(2). D-II/CH/1, G.I.D.C. Estate, Dahej, Tal. Vagra, Dist. Baruch,
Gujarat-392130
Gujarat

(2).

(3). Plot No 905 , GIDC Estate, Anklehswar
Gujarat

(3).

(4).

(4). 226/1,226/2 & 227/1 G.I.D.C Estate, Panoli,
Gujarat-394116,India
Gujarat

(5).

(5). 228, G.I.D.C. Estate, Panoli,Anklehswar, Bharuch - 394 116,
Gujarat State
Gujarat

Customer Care Details

»—« ± μ

Safety Pictograms:



Big Label

(> 250 mL Primary and Secondary Package Label)

Read leaflet before use:

Common Name:

Tebuconazole 6.7% Captan 26.9% w/w SC

BRAND NAME: -

Crops and Pest Groups: Tebuconazole 6.7% + Captan 26.9% w/w SC is a combination of fungicide which is used for the control of Powdery mildew and Anthracnose disease of Chilli, Powdery mildew, Alternaria leaf spot/ blight and scab of Apple.

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For agriculture use.

Re-entry:

24 hours.

Chemical Composition :

Content % w/w

Captan a.i.	active ingredient	26.9 % w/w
Tebuconazole a.i	active ingredient	6.7 % w/w
Polyarylphenyl ether phosphate	other ingredient	2.1 % w/w
Propylene glycol	other ingredient	5.9 % w/w
Xanthan Gum	other ingredient	0.25 % w/w
Water	other ingredient	q.s % w/w
Modified-dimethyl polysiloxane	other ingredient	0.6 % w/w
Sodium Naphthalene formaldehyde condensate	other ingredient	1 % w/w

Total:100.00%

Tebuconazole 6.7% Captan 26.9% w/w SC

Fungicide



Keep out of reach of children.

Keep in cool, dry place away from heat and open flame.

Regn. No. _____

Mfg. Lic. No. _____

Batch No. _____

Mfg Date _____

Exp. Date _____

Net Content / Volume _____

Max. Retail Price _____

3 TM - TSSISS

Safety Precautions:

Premises Address

(1). S.No. 180/6, & 180/7, Khazipally (V), Jinnaram (M), Dist-Medak, Telangana-502 319

(2). D-II/CH/1, G.I.D.C. Estate, Dahej, Tal. Vagra, Dist. Baruch, Gujarat-392130 Gujarat

(3). Plot No 905 , GIDC Estate, Anklehswar Gujarat

(4). 226/1,226/2 & 227/1 G.I.D.C Estate, Panoli, Gujarat-394116,India Gujarat

(5). 228, G.I.D.C. Estate, Panoli,Anklehswar, Bharuch - 394 116, Gujarat State Gujarat

© 2011 ©

(1).

(2).

(3).

(4).

(5).

Customer Care Details

» « ± μ

Safety Pictograms:



Tebuconazole 6.7% Captan 26.9% w/w SC

(Fungicide)

Tebuconazole 6.7% + Captan 26.9% w/w SC is a combination of fungicide which is used for the control of Powdery mildew and Anthracnose disease of Chilli, Powdery mildew, Alternaria leaf spot/ blight and scab of Apple. Tebuconazole 6.7% + Captan 26.9% w/w SC is a combination of fungicide which is used for the control of Powdery mildew and Anthracnose disease of Chilli, Powdery mildew, Alternaria leaf spot/ blight and scab of Apple.

Chemical Composition:	Ingredient	%W/W
Captan a.i.	active ingredient	26.9 % w/w
Modified-dimethyl polysiloxane	other ingredient	0.6 % w/w
Polyarylphenyl ether phosphate	other ingredient	2.1 % w/w
Propylene glycol	other ingredient	5.9 % w/w
Sodium Naphthalene formaldehyde condensate	other ingredient	1 % w/w
Tebuconazole a.i	active ingredient	6.7 % w/w
Water	other ingredient	q.s % w/w
Xanthan Gum	other ingredient	0.25 % w/w
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water L	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI Gram	Formulation ML					
Chilli	Powdery Mildew and Anthracnose	80+320	1000	500	Vegetative stage and at the time of appearance of disease.	24 hours	5	2-3 sprays based on infestation.
Apple	Powdery mildew, Alternaria leaf spot/blight, Scab	124 + 496 or 0.02%+0.08%	1550	10 L / tree	Vegetative stage and at the time of appearance of disease.	24 hours	10	2-3 sprays based on infestation.

Note:-

Good Practices relating to pesticide application techniques

Direction of Use

The required quantity of ready mix chemical Tebuconazole 6.7%+Captan 26.9% w/w SC need to be diluted in water and sprayed by using knapsack sprayer / foot sprayer fitted with hollow cone nozzle with spray volume @ 500 l/ha for Chilli and 10 L/tree for Apple.

Weather Condition(in general):

Should be used during favorable weather conditions and should not be used during extreme weather conditions.

Soil & Water(where ever applicable):

Light to heavy soil. Clean water to be used for spray.

Crop Stage for application:

Vegetative stage and at the time of appearance of disease.

Application Technique

Foliar application with the help of Knapsack sprayer / foot sprayer.

Equipment Type:

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Knapsack sprayer / foot sprayer.

NozzleType:

Hollow cone nozzle.

Dosage:

Please refer recommendation table.

Instruction for Mixing:

Please see " direction of use".

Re-entry Period

24 hours

Manner and Time of Application

Should be used preferably in early morning or evening hours.

Application Timing

To be applied preventively and to be repeated at 15 days interval depending upon prevailing weather conditions and disease pressure.

Frequency

2-3 sprays based on infestation.

Stages during which pesticides should not be applied on any crop :

Not to be applied during crop maturity and just before crop harvest.

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

not to be used on crops other than mentioned on label & leaflets. 2. it should not be used in the areas where pisciculture / aquaculture is practiced and honey bee.

Symptoms Of Poisoning

Weakness, fatigue, Dyspnea, Blurred vision, Irritation of eyes, Skin and Upper respiratory system, Dermatitis, Skin sensitization, Diarrhoea, vomiting and Conjunctivitis, nausea, headache.

First Aid

If ingested: If person is conscious and alert, rinse mouth with water and give 1 or 2 glass of water to drink. Never give anything by mouth to an unconscious person. Call the doctor. Do not induced vomiting until and unless desired by attending Physician. If Inhaled: Take the patient to fresh air and take care of the body temperature. If patient is not breathing or breathing with difficulty, give artificial respiration, preferably with mouth to mouth breathing. Consult a physician. In case of skin contact: Wash thoroughly the affected part of body with soap and plenty of water including hairs and under fingernails. If pain, redness or irritation persists, consult a doctor. All the contaminated clothes should segregated and thoroughly washed with soap and water separately before use. In case of Eye contact: Rinse gently with plenty of water at least for 15 minutes. If pain, irritation, redness or photophobia persist. Consult eye specialist.

Antidote

No specific Antidote, Treat symptomatically.

Phytotoxicity

The product is not phytotoxic when used as per recommendation.

Storage Conditions

1. The package containing pesticide shall be stored in separate rooms or premises away from storing other articles or shall be in separate almirahs under lock and key depending upon the quantity and nature of the pesticide. 2. The rooms or premises means for storing pesticides shall be well built, dry, well-lit and ventilated and sufficient dimension to prevent environmental contamination with vapour of pesticides.

Safety Precaution:

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

During spraying wear suitable respiratory equipment. Wash clothing before re-using. Wash hands thoroughly after handling. Wear protective clothing like suitable gloves, chemical resistant boots, Chemical goggles or safety glasses. Spray only in the wind blowing direction only not against so as to prevent the settling drifts on body. Do not eat, drink, chew or smoke or handle food while spraying. Avoid contact with skin and eyes.

Cautionary Statement:

1. Not to be used on crops other than mentioned on label & leaflets.

2. Not should not be used in the areas where pisciculture / aquaculture, honey bee is practiced.

Disposal of used containers, washing from equipment and surplus material :

1.It shall be the duty of manufacturers, formulators of insecticides and operators to dispose packages or surplus materials and washing in a safe manner so as to prevent environmental or water pollution. 2. The used packages shall not be left outside to prevent their re-use. 3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

226/1,226/2 & 227/1 G.I.D.C Estate, Panoli, Gujarat-394116,India
Telangana

228, G.I.D.C. Estate, Panoli,Ankleshwar, Bharuch - 394 116,
Gujarat State
Telangana

D-II/CH/1, G.I.D.C. Estate, Dahej, Tal. Vagra, Dist. Baruch,
Gujarat-392130
Telangana

Plot No 905 , GIDC Estate, Ankleshwar
Telangana

S.No. 180/6, & 180/7, Khazipally (V), Jinnaram (M), Dist-Medak,
Telangana- 502 319
Telangana

Manufactured by

ADAMA India Pvt. Ltd.
Plot No: DS-13, IKP Knowledge Park, Sy. No. 542/2, Genome
Valley, Turkapally, Shameerpet, Ranga Reddy District Hyderabad
Telangana 500078

Customer Care Details:

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet