

Primary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888

BRAND NAME: -

Chemical Composition :

Content % w/w

Carboxy methyl cellulose	1.0% w/w
Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888	1.0% wp
Talc Powder	98.0% w/w

Total: 100.00%

Please refer leaflet for details

Antidote

Treat Symptomatically.

Manufactured by

TStanes and Company Limited,
8/23-24, Race Course Road, Coimbatore- 641 018 Coimbatore
Coimbatore Tamil Nadu

Premises Address

(1). 8/23-24, Race Course Road, Coimbatore- 641 018
Tamil Nadu

(2). 502/1, 503/1A, Pilladam Road, Okkillipalayam,
Othakalmandabam, Coimbatore - 641032

(3). 30/31, Railway Station Road, Thudiyalur, Coimbatore 641034

Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888

o—|² |>’j x-²’ \$!#’ ’\$!’@!’ (IIHR TH-2 Strain Accession No. ITCC N0 6888

Fungicide + Insecticide



o—|² |>’j x-²’ \$!#’ ’\$!’@!’ (IIHR TH-2 Strain Accession No. ITCC N0 6888

Keep in cool, dry place away from heat and open flame.

Regn. No.	’ j !’-!	_____
Mfg. Lic. No.	2 - ₃ !’ μ !’-!	_____
Batch No.	° œ’-!	_____
Mfg Date	2 - ₃ !’ @!	_____
Exp. Date	¿ ñ²’ » -’-!’ @’-’	_____
Net Content / Volume	, ’ ± ’ž ñ ³ @-	_____
Max. Retail Price	m ñ—@ ² -’ » ’ ² ý ³	_____
Customercare		_____

q@³ TM’ o’ ’@’ μ’ @’ —’ @É-

o—|² |>’j x-²’ \$!#’ ’\$!’@!’ (IIHR TH-2 Strain Accession No. ITCC N0 6888

—Á

’ o³ —’-’ ° œ’- -

—° a° o’² a oμ’ 1.0% w/w

° y³ μ°

o—|² |>’j x-²’ 2 x 10^6 / 2’ ò³ -! 1.0% wp

\$!#’ ’\$!’@!’ (IIHR TH-2 Strain Accession No. ITCC N0 6888

μ —’ @q!’ 98.0% w/w

3 TM’ - TSS!SS’

μ ¼” —’ q@œ’-’-’ Q

qñ@«—

μ!’ ñò° ’wò!’-ú@-’ μ² ñ!ž

8/23-24!’ ° o’-’ ° |’ j ž —³ 2° ñ’ ž’ 641 018’ —³ 2° ñ’

@’ o’ -’-’ @@-

(1). 8/23-24. ??? ????? ???, ????????? - 641 018

(2). 502/1, 503/1A, Pilladam Road, Okkillipalayam, Othakalmandabam, Coimbatore - 641032

(3). 30/31, Railway Station Road, Thudiyalur, Coimbatore 641034

Secondary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888

BRAND NAME: -

Crops and Pest Groups: Trichoderma harzianum 1% WP is a bio-fungicide which effectively control Root-knot nematodes and Wilt disease in Tomato and Okra crops.

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For agricultural use

Re-entry:

12 hours.

Chemical Composition :

Content % w/w

Carboxy methyl cellulose		1.0% w/w
Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888	2x10 ⁶ /gm.min	1.0% wp
Talc Powder		98.0% w/w

Total: 100.00%

Safety Precautions, Symptom of Poisoning and First Aid :Please refer leaflet for details.

Antidote Statement:

Treat Symptomatically.

Cautionary statement:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888

o—|² |»¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888
Fungicide + Insecticide



o—|² |»¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

Keep in cool, dry place away from heat and open flame.

Regn. No. _____
Mfg. Lic. No. _____
Batch No. _____
Mfg Date _____
Exp. Date _____
Net Content / Volume _____

q®³ TM^o | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

o—|² |»¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

—A

—1¹ 3 TM^o | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

\$% \$ α

1^o 3 —^o | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

—^o â^o | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

° y³ μ °

o—|² |»¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

\$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

α —¹ | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

3 TM^o | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

μ ¼^o | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

\$! μ ° μ | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

q «³ —¹ | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

@² | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

@ ° o | »¹ | x-² | \$!#[~] | \$! | @! | (IIHR TH-2 Strain Accession No. ITCC NO 6888

Manufactured by

TStanes and Company Limited,
8/23-24, Race Course Road, Coimbatore- 641 018 Coimbatore
Coimbatore Tamil Nadu

Premises Address

(1). 8/23-24, Race Course Road, Coimbatore- 641 018
Tamil Nadu

(2). 502/1, 503/1A, Pilladam Road, Okkillipalayam,
Othakalmandabam, Coimbatore - 641032

(3). 30/31, Railway Station Road, Thudiyalur, Coimbatore 641034

Max. Retail Price

qñ®«—

8/23-24. 641 018

(2). 502/1, 503/1A, Pilladam Road, Okkillipalayam, Othakalmandabam, Coimbatore - 641032

(3). 30/31, Railway Station Road, Thudiyalur, Coimbatore 641034

Customer Care Details

Safety Pictograms:



Big Label

(> 250 mL Primary and Secondary Package Label)

Read leaflet before use:

Common Name:

Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888

BRAND NAME: -

Crops and Pest Groups: Trichoderma harzianum 1% WP is a bio-fungicide which effectively control Root-knot nematodes and Wilt disease in Tomato and Okra crops.

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For agricultural use

Re-entry:

12 hours.

Chemical Composition :

	Content % w/w
Carboxy methyl cellulose	1.0% w/w
Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888	1.0% wp
Talc Powder	98.0% w/w
Total:100.00%	

Safety Precautions:

Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888

o—|² l»'j x-² '\$!#~ 's!'@!' (IIHR TH-2 Strain Accession No. ITCC N0 6888

Fungicide + Insecticide



o—|² l»'j x-² '\$!#~ 's!'@!' (IIHR TH-2 Strain Accession No. ITCC N0 6888

Keep in cool, dry place away from heat and open flame.

- Regn. No. _____
- Mfg. Lic. No. _____
- Batch No. _____
- Mfg Date _____
- Exp. Date _____
- Net Content / Volume _____
- Max. Retail Price _____

g®³ TM ° '®µ'® —'®E-

o—|² l»'j x-² '\$!#~ 's!'@!' (IIHR TH-2 Strain Accession No. ITCC N0 6888

—Á

..... A

—1' 3 TM —' µw

\$% § π

..... 2 ± ±

—° á° ' 2ª oµ' 1.0% w/w

° y³ µ °

o—|² l»'j x-² ' 2 x 10^6 / ² · 0³ —! 1.0% wp

\$!#~ 's!'@!' (IIHR TH-2 Strain Accession No. ITCC N0 6888

π —'®q! ' 98.0% w/w

..... 3 TM - TSSISS'

Customer Care Details

» « ± µ · · ° °

Safety Pictograms:



fl' ' ' « - , -f' -m- , -fl

2 x 10⁶ gm.min

o-|² l»'j x-² \$!#~ 'š!@!' (IIHR TH-2 Strain Accession No. ITCC N0 6888

° 3 - - ° (œ- ' -

—° â° ' 2 ^a oμ'° y³ μ°	1.0% w/w
±-i'@q '	98.0% w/w
o- ² l»'j x-² \$!#~ 'š!@!' (IIHR TH-2 2 x 10 ⁶ / ²'ô³-!	1.0% wp
Strain Accession No. ITCC N0 6888	

3 TM - TSS!SS~ '± '± '± '±

° μ'	—m'—-2	©'»â³'12		° μ'm' a	@- '—μ	@ l' ° μ	n .
		° l' ©ñ	° 'œ-				
±² ±'	2 μ ô ^a ' ° 2' ff μ o TML' - o—TML ± fl q—m' ' TM' fê.3 ° '3 2' . { â° @'2' m' 'w' @' μ o—®' ° —fl	° l' ©ñ	w— 'l' !' ° j' —' %#' 2' o— ² l »'j x-² \$!#~ 'š!@!' ° ° j' @œ' —' ° a' . —° l' ° j' —' i' . — —m- , —' ° (#' 2' ° ° . TM' 2 !' —' « ' ° ° ° q@œ' ©' —' Q ñ³ ' ®' . ° ° @»μ' (' —' !' !' o— ² l»'j x-² \$!#~ 'š!@!' ° ° ° . - @k' (' m- ' TM° ° ° —' « ' —' ©' »â³'12 —' « ' ° ° ± 2' 2' 3 TM' —'	ñ³ ' ®' ° ° @»μ	\$% š ±	j' . —' —m- , —' — μwn . 3 —' —»	w—' ° ° ° j' @œ' . ' w—' ° ° ° ± 2 q@œ' —' μw ° . p' ° ° @»μ
±	2 μ ô ^a ' ° 2' ff μ o TML' - o—TML ± fl q—m' ' TM' fê.3 ° '3 2' . { â° @'2' m' 'w' @' . ° - â² fl	° l' ©ñ	w— 'l' !' ° j' —' %#' 2' o— ² l »'j x-² \$!#~ 'š!@!' ° ° j' @œ' —' Q ° . p' ° ° @»μ' (' —' !' !' o— ² l»'j x-² \$!#~ 'š!@!' ° ° ° . - @k' (' m- ' TM° ° ° —' « ' —' ©' »â³'12 —' « ' ° ° ± 2' 2' 3 TM' —' Q	° . p' ° ° @»μ	\$% š ±	j' . —' —m- , —' — μwn . 3 —' —»	w—' ° ° ° j' @œ' . ' w—' ° ° ° ± 2 q@œ' —' μw ° . p' ° ° @»μ

Note:-

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

502/1, 503/1w äµ «² º º | Ž { —µ äµ ³ ² Ž { º —µ² —| º ² Ž —³ úº @ º º >
641032

8/23-24! º º —º | º | Ž —³ ² º º º º 641 018

»—« ± µ º º º º

º º ¼ º º º µ º



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Trichoderma harzianum 1.0% WP (IIHR TH-2 Strain Ac.No. ITCC-6888

(Fungicide + Insecticide)

2 x 10⁶ gm.min

Trichoderma harzianum 1% WP is a bio-fungicide which effectively control Root-knot nematodes and Wilt disease in Tomato and Okra crops.

Chemical Composition:	Ingredient	%W/W
	Carboxy methyl cellulose	1.0% w/w
	Talc Powder	98.0% w/w
	Trichoderma harzianum 1.0% WP (IIHR TH-2 2x10 ⁶ /gm.min Strain Ac.No. ITCC-6888	1.0% wp
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI	Formulation					
Tomato	Root-knot nematodes (Meloidogyne incognita) Wilt (Fusarium oxysporum f. sp. lycopersici)	-	Treat the seeds with Trichoderma harzianum 1.0% WP @ 20 g/kg of seeds; treat the nursery beds with the biopesticide @ 50 g/sq m and apply 5 tons FYM enriched* with 5 kg of Trichoderma harzianum 1.0% WP/ hectare to the soil before transplanting	-	Before transplanting	12 hours	Not applicable for bio-pesticides	one time as seed treatment and once as soil application
Okra	Root-knot nematodes (Meloidogyne incognita) Wilt (Fusarium oxysporum f. sp. vasiinfectum)	-	Treat the seeds with Trichoderma harzianum 1.0% WP @ 20 g/kg of seeds and apply 5 tons FYM enriched* with 5 kg of Trichoderma harzianum 1.0% WP/ hectare to the soil before sowing.	-	Before sowing	12 hours	Not applicable for bio-pesticides	one time as seed treatment and once as soil application

Note:-

Good Practices relating to pesticide application techniques

Direction of Use

Mix required quantity of Trichoderma harzianum 1.0% WP with seed and ensure uniform coating. Enrich one kg formulation with one ton farm yard manure or compost and left under shade for 15 days maintaining optimum moisture. Seed treatment using drums or by hand wearing rubber gloves. Broadcast by hand wearing rubber gloves. Do not apply Trichoderma harzianum 1.0% WP on chemically treated seeds.

Weather Condition(in general):

Should be used during favorable weather conditions and should not be used during extreme weather conditions.

Soil & Water(where ever applicable):

Light to heavy soil. Clean water to be used.

Crop Stage for application:

Before transplanting

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

Application Technique

Seed treatment and soil treatment

Equipment Type:

Seed treatment drums or by hand wearing rubber gloves. Broadcasting by hand wearing rubber gloves.

NozzleType:

Not applicable

Dosage:

Tomato: Treat the seeds with Trichoderma harzianum 1.0% WP @ 20 g/kg of seeds; treat the nursery beds with the biopesticide @ 50 g/sq m and apply 5 tons FYM enriched* with 5 kg of Trichoderma harzianum 1.0% WP/ hectare to the soil before transplanting

Okra: Treat the seeds with Trichoderma harzianum 1.0% WP @ 20 g/kg of seeds and apply 5 tons FYM enriched* with 5 kg of Trichoderma harzianum 1.0% WP/ hectare to the soil before sowing.

Instruction for Mixing:

Do not apply Trichoderma harzianum 1.0%WP on chemically treated seeds.

Re-entry Period

12 hours

Manner and Time of Application

Tomato: before transplanting.

Okra: before sowing.

Application Timing

Should be used preferably in early morning or evening hours

Frequency

one time as seed treatment and once as soil application

Stages during which pesticides should not be applied on any crop :

Not applicable as the product is applied as seed treatment/soil treatment before sowing/transplanting.

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

Symptoms Of Poisoning

Headache, Nausea, Vomiting, irritation & allergy may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to 15 minutes. 4. Remove the patient into fresh air.

Antidote

Treat Symptomatically.

Phytotoxicity

Non phytotoxic at recommended dosage

Storage Conditions

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

1. The packages containing the Bio-fungicide shall be stored in separate room or premises away from the rooms or premises used for storing other articles particularly articles of food or shall be kept in separate almirahs under lock and key depending upon the quantity and nature of the Bio-fungicide.
2. The rooms or premises meant for storing the Bio-fungicide shall be well built, dry, well lit and ventilated and of sufficient dimension to avoid contamination with its vapours.

Safety Precaution:

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothings which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

Cautionary Statement:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufacturers, formulators of insecticides and operators to dispose packages or surplus materials and washing in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

30/31, Railway Station Road, Thudiyalur, Coimbatore 641034
Tamil Nadu

502/1, 503/1A, Pilladam Road, Okkillipalayam, Othakalmandabam,
Coimbatore - 641032
Tamil Nadu

8/23-24, Race Course Road, Coimbatore- 641 018
Tamil Nadu

Manufactured by

TStanes and Company Limited,
8/23-24, Race Course Road, Coimbatore- 641 018 Coimbatore
Tamil Nadu

Customer Care Details:

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet