

Primary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

BRAND NAME: -

Chemical Composition :

Content % w/w

Strain-IIHR-TV-5, Acc.No.ITCC-6889	(CFU count 1 x 10 ⁶ /gm min.)	1.50 % w/w
Carboxy Methyl Cellulose		1.5 % w/w
Talc Powder		97.0% w/w
total	Total	100.00% w/w

Total: 100.00%

Please refer leaflet for details

Antidote

Treat Symptomatically.

Manufactured by

TROPICAL AGROSYSTEMS (INDIA) PVT. LTD.
72, MARSHALL'S ROAD, EGMORE CHENNAI-600008 CHENNAI
Chennai Tamil Nadu 600008

Premises Address

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Strain-IIHR-TV-5, Acc.No.ITCC-6889

Contact Fungicide

சாத்திரம் - 6889



சாத்திரம் - 6889

Keep in cool, dry place away from heat and open flame.

Regn. No.	சாத்திரம் - 6889
Mfg. Lic. No.	2-31-μ-1-1
Batch No.	°œ-!
Mfg Date	2-31-©!
Exp. Date	2-31-©!
Net Content / Volume	1.5 ± 0.1 g
Max. Retail Price	₹ 100.00
Customercare	600008

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Strain-IIHR-TV-5, Acc.No.ITCC-6889

Contact Fungicide

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Strain-IIHR-TV-5, Acc.No.ITCC-6889	(CFU count 1 x 10 ⁶ /gm min.)	1.50 % w/w
Carboxy Methyl Cellulose		1.5 % w/w
Talc Powder		97.0% w/w
total	Total	100.00% w/w

Total: 100.00%

(1). 530/2-B, Vanagaram Road, Ambattur, Chennai-600058,
Tamilnadu, India
Tamil Nadu

(2). 80/3/3, Arya Nagar & 34-36,Rural Industrial Area, Loni, District-
Ghaziabad, Uttar Pradesh, India.
Uttar Pradesh

(3). Plot No. A/74-1 & A/74-2, U.P.S.I.D.C, Sikandrabad,
Bulandshahar, Uttar Pradesh, India.
Uttar Pradesh

(1). 530/2-??, ????????, ????????, ???????-600058, ????????,
????

(2). 80/3/3, ?????? ?? 34-36, ????, ?????????????? ??, ????,
?????????- ??????????, ??????????, ????

(3). ?????? ?? ?/74-1 ?? ?/74-2, ???.???.???.??.?, ???????????,
?????????, ??????????, ????

Customer Care Details

»—« ± μ

Safety Pictograms:



Secondary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

BRAND NAME: -

Crops and Pest Groups: Trichoderma viride 1.5% WP is a bio-fungicide which effectively control Root-knot nematodes and Wilt disease in Tomato and Okra crops.

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For Agriculture use

Re-entry:

12 hours

Chemical Composition :

Content % w/w

Strain-IIHR-TV-5, Acc.No.ITCC-6889	(CFU count 1 x 10 ⁶ /gm min.)	1.50 % w/w
Carboxy Methyl Cellulose		1.5 % w/w
Talc Powder		97.0% w/w
total	Total	100.00% w/w

Total: 100.00%

Safety Precautions, Symptom of Poisoning and First Aid :Please refer leaflet for details.

Antidote Statement:

Treat Symptomatically.

Cautionary statement:

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889
Contact Fungicide

300 gms



300 gms

Keep in cool, dry place away from heat and open flame.

Regn. No.		_____
Mfg. Lic. No.		_____
Batch No.		_____
Mfg Date		_____
Exp. Date		_____
Net Content / Volume		_____

300 gms

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

300 gms

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

For Agriculture use

12 hours

1.50 % w/w

1.5 % w/w

97.0% w/w

total Total 100.00% w/w

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

300 gms

Safety Precautions, Symptom of Poisoning and First Aid :Please refer leaflet for details.

Antidote Statement:

Treat Symptomatically.

Cautionary statement:

Big Label

(> 250 mL Primary and Secondary Package Label)

Read leaflet before use:

Common Name:

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

BRAND NAME: -

Crops and Pest Groups: Trichoderma viride 1.5% WP is a bio-fungicide which effectively control Root-knot nematodes and Wilt disease in Tomato and Okra crops.

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For Agriculture use

Re-entry:

12 hours

Chemical Composition :

	Content % w/w
Strain-IIHR-TV-5, (CFU count 1 x 10 ⁶ /gm min.)	1.50 % w/w
Carboxy Methyl Cellulose	1.5 % w/w
Talc Powder	97.0% w/w
total	Total 100.00% w/w

Total:100.00%

Safety Precautions:

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

Strain-IIHR-TV-5, Acc.No.ITCC-6889

Contact Fungicide

® 3 1 1 « - , -



o - | ' 2 ' ' | ' \$! (' ' \$ μ - , μ ' œ ' l ' 4 ' - - - - - 6 Strain-IIHR-TV-5, Acc.No.ITCC-6889

Keep in cool, dry place away from heat and open flame.

Regn. No.	_____
Mfg. Lic. No.	_____
Batch No.	_____
Mfg Date	_____
Exp. Date	_____
Net Content / Volume	_____
Max. Retail Price	_____

q®³ TM ° ' ® μ ' ® - ' ® E-

o - | ' 2 ' ' | ' \$! (' ' \$ μ - , μ ' œ ' l ' 4 ' - - - - - 6 Strain-IIHR-TV-5, Acc.No.ITCC-6889

— Á

..... A

— 1 ' q®³ TM ' — μ w!

\$% \$ π

1 ° 3 - - - ° œ - 2 ' ± ±

o - | ' 2 ' ' | ' \$! # ' ' 4 - μ - ' - 2 l ' w - - - fl 1.50 % w/w

\$ μ - , μ ' œ ' l ' 4 ' - - - ° l ' ! 3 ' fl - 1 x 10⁶ /

— - - - - 6 Strain-IIHR-TV-5, Acc.No.ITCC-6889

— ° á ° ' 2 - - o μ ' 1.5 % w/w

° ý³ μ °

π ý - ' ® q l ' 97.0% w/w

— μ - - - - - μ 100.00% w/w

3 TM - TSSISS

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothings which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

Symptom of Poisoning:

Headache, Nausea, Vomitting, irritation & allergy may occur.

First Aid:

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to15 minutes. 4. Remove the patient into fresh air.

Antidote

Treat Symptomatically.

Cautionary statements:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Manufactured by

TROPICAL AGROSYSTEMS (INDIA) PVT. LTD.
72, MARSHALL'S ROAD, EGMORE CHENNAI-600008 CHENNAI
Chennai Tamil Nadu 600008

Premises Address

(1). 530/2-B, Vanagaram Road, Ambattur, Chennai-600058,
Tamilnadu, India
Tamil Nadu

(2). 80/3/3, Arya Nagar & 34-36,Rural Industrial Area, Loni, District-
Ghaziabad, Uttar Pradesh, India.
Uttar Pradesh

(3). Plot No. A/74-1 & A/74-2, U.P.S.I.D.C, Sikandrabad,
Bulandshahar, Uttar Pradesh, India.
Uttar Pradesh

It is a document with illegible text, possibly a scanned image of a document where the text is too small or blurry to read. The text appears to be a list of items or a table with multiple columns.

It is a document with illegible text, possibly a scanned image of a document where the text is too small or blurry to read. The text appears to be a list of items or a table with multiple columns.

It is a document with illegible text, possibly a scanned image of a document where the text is too small or blurry to read. The text appears to be a list of items or a table with multiple columns.

It is a document with illegible text, possibly a scanned image of a document where the text is too small or blurry to read. The text appears to be a list of items or a table with multiple columns.

(1) 530/2-??, ??????? ??, ???????, ???????-600058, ???????,
????

(2) 80/3/3, ????? ?? ? 34-36, ??? ???????????? ???, ???,
????????- ?????????, ?????????, ????

(3) ????? ? ?/74-1 ? ?/74-2, ???.??.??.??.??.?, ?????????,
????????, ?????????, ????

Customer Care Details

» « ± μ ∙ ∙ ∙ »

Safety Pictograms:



fi 3 2 1

2 x 10 gm.min

Q

3 2 1

2 3 1	1.5 % w/w
4 2 1 / 1x10^6	1.50 % w/w
6 Strain-IIHR-TV-5, Acc.No.ITCC-6889	2 3 1
2 3 1	97.0% w/w
2 3 1	100.00% w/w
3 2 1 TSSISS 2 1	

	2 3 1	2 3 1			2 3 1	2 3 1	n	
		2 3 1	2 3 1					
2 3 1	2 3 1 o- f-3 j	2 3 1	T- 2 3 1 2 3 1 US o- s! o- s! 2 o Q o- TIX s! - o- TIX s! 2 2 o Q o- TIX s! - ! o- X o 3	2 3 1	2 3 1 Q	\$% s	-	2 3 1 2 3 1 w
2 3 1	2 3 1 o- f-3 j	2 3 1	T- 2 3 1 2 3 1 US o- s! o- s! 2 o Q o- TIX s! - ! o- X o 3	2 3 1	2 3 1 p	\$% s	-	2 3 1 2 3 1 w

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Note: 1 →μ 2 ° 1œ 2 ° 0—|2 | . 1° —:\$(2 ° 3 2 ° » © ' »

q®3 TM — μw —« .

μ2 μ—T —μ 2 ° 1° —2 ° —US 2 ° 0—|2 | . 1° TIX ° Š!œ 1° ° j q®œ —° © 1° —° |° —° 0—|2 | . 1° TIX ° Š!œ 1° —2 ° XS 2 ° TM 2 μ —« ° ° j q®œ —° © ° ° μ ± 2 ° —° 0—|2 | . 1° TIX ° Š!œ 1° fx' —° !"»â μ3 fl ° ° . « a ° TM ° —° « ' E —' X' μ —"»â μ3 —° « ° ° 3 TM —!

±0| —T —μ 2 ° 1° —2 ° —US 2 ° 0—|2 | . 1° TIX ° Š!œ 1° ° j ®œ —° © 1° . p ° ° μ ± 2 ° —° 0—|2 | . 1° TIX ° Š!œ 1° fx' —° !"»â μ3 fl ° ° . « a ° TM ° —° « ' E —' X' μ —"»â μ3 —° « ° ° 3 TM —!

2 ° 2 ° —° a ©

2 ° 2 ° —° m—μ ° ® ° a ©3 —° « ° —° q®3 TM —3 ° j —° œ »w ° a ° ©—μ ° « , { —° « ° —° o ° ©2 μ ° —° Q

2 ° μ | ° ° —°

»μ —° ° ± j ° 2 μ μ ° © —° μ w Q ° Ó ° ® —° —° o ° ©2 μ ° —° Q

° μ ° m ° a

© ° ® ° ° ° ® μ Q

n . « ° —° © —° —°

° j ° q®œ ° | ° ° 2 ° « ° m—° 3 TM

° j ° q®œ ° ú ° 3 ° ° ° —° « ° —° ° ± 2 ° q®œ ° —° μ w ° ° ° —° « ° —° ° ® —° —° » a ° ° ° Ÿ | —3

μ ° TM —° » ° °

Please refer recommendation table

—®3 ° ° q®3 TM —° —° « ° ° ° « ° —° Q

° ° 3 —° —° q®œ ° © ° ° j ° ° ® ° —° μ ° TM 3 Q

\$% Š μ

—®3 ° ° q®3 TM —° —° « ° ° ° « ° —° Q

n . « ° —° —° ° 2 3

° ° » ° 3 ° ° 2 ° —° ° 2 3 ° 2 ° ° ©2 μ ° —° 3 ° j —° œ »w!

w—° ° ° ° j ° ®œ ° | ° ° w—° ° ° ± 2 q®œ ° —° μ w

μ ° TM —° » ° » ° â 3 —° o ° ° qñ® « ° —° 3 TM ° ° j ° ®œ ° ± 2 q®œ ° 2 ° ° . p ° ° © ° ® ° ° ° ® μ ° —° 3 ° j ° © ° » Q

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet!

80/3/3Ž n 3 | -™ | 34-36Ž 'μ' o| 3 μ' x ' n Ž μ - Ž | ° μ ' - μ ž
™ j 3 ° «Ž q©' «, Ž ± ' ©!

öμ μ' -! ' x"74-1' | ' ' x"74-2Ž 3! @! w! In p! | !° ! Ž ° -« ' ° «Ž ° μ«, »' Ž
q©' «, Ž ± ' ©

»-«' ± μ' . . ' -

° ' ¼ ' œ ' μ ' ®-



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Trichoderma Viride 1.5% WP Strain-IIHR-TV-5, Acc.No.ITCC-6889

(Contact Fungicide)

2 x 10⁶ gm.min

Trichoderma viride 1.5% WP is a bio-fungicide which effectively control Root-knot nematodes and Wilt disease in Tomato and Okra crops.

Chemical Composition:	Ingredient	%W/W
	Carboxy Methyl Cellulose	1.5 % w/w
	Strain-IIHR-TV-5, Acc.No.ITCC-6889 (CFU count 1 x 10 ⁶ /gm min.)	1.50 % w/w
	Talc Powder	97.0% w/w
	total	Total 100.00% w/w
		Total: 100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI	Formulation					
Tomato	Root-knot nematodes (Meloidogyne incognita) Bacterial Wilt (Ralstonia solanacearum)	-	Treat the seeds with Trichoderma viride 1.5% WP @ 20 gm/kg of seeds & nursery beds with the Trichoderma viride 1.5% WP @ 50 gm/sq.m. and also apply Trichoderma viride 1.5% WP (@ 5 kg/hectare) enriched FYM* @ 5 tons/hectare to the soil before transpla	-	Before transplanting	12 hours	-	one time as seed treatment and once as soil application
Okra	Root-knot nematodes (Meloidogyne incognita) Wilt (Fusarium oxysporum)	-	Treat the seeds with Trichoderma viride 1.5% WP @ 20 gm/kg of seeds and Trichoderma viride 1.5% WP@ 5 kg/hectare enriched FYM* @ 5 tons/hectare to the soil before sowing.	-	Seed treatment before crop sowing	12 hours	-	one time as seed treatment and once as soil application

Note:- 1 kg of formulation contains 15 gm biomass of Trichoderma viride

Good Practices relating to pesticide application techniques

Direction of Use

Tomato: Treat the seeds with Trichoderma viride 1.5% WP @ 20 gm/kg of seeds & nursery beds with the Trichoderma viride 1.5% WP @ 50 gm/sq.m. and also apply Trichoderma viride 1.5% WP (@ 5 kg/hectare) enriched FYM* @ 5 tons/hectare to the soil before transplanting.

Okra: Treat the seeds with Trichoderma viride 1.5% WP @ 20 gm/kg of seeds and Trichoderma viride 1.5% WP@ 5 kg/hectare enriched FYM* @ 5 tons/hectare to the soil before sowing.

Weather Condition(in general):

Should be used during favorable weather conditions and should not be used during extreme weather conditions.

Soil & Water(where ever applicable):

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Light to heavy soil. Clean water to be used.

Crop Stage for application:

Before transplanting

Application Technique

seed treatment and soil application

Equipment Type:

Seed treatment drums or by hand wearing rubber gloves. Broadcasting by hand wearing rubber gloves.

NozzleType:

Not applicable

Dosage:

Please refer recommendation table

Instruction for Mixing:

Please see "direction of use".
Do not apply on chemically treated seeds.

Re-entry Period

12 hours

Manner and Time of Application

Please see "direction of use"

Application Timing

Should be used preferably in early morning or evening hours.

Frequency

one time as seed treatment and once as soil application

Stages during which pesticides should not be applied on any crop :

Not applicable as the product is applied as seed treatment/soil treatment before sowing/transplanting.

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

Symptoms Of Poisoning

Headache, Nausea, Vomitting, irritation & allergy may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to15 minutes. 4. Remove the patient into fresh air.

Antidote

Treat Symptomatically.

Phytotoxicity

Not Phytotoxic at recommended dosage.

Caution:Í Not to Be Used on Crops Other Than Specified on This Label/LeafletÍ "

Storage Conditions

- 1) The package containing Biological pesticide shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles or shall be kept in separate almirahs under lock & key depending upon quantity & nature of the Biological pesticide.
- 2) The rooms or premises meant for storing Biological pesticide shall be well built, dry, well-lit and ventilated and be of sufficient dimension to prevent environmental contamination.
- 3) The product should be stored below 45°C temperature away from direct sunlight.
- 4) Store separately from bactericidal and chemical pesticides.

Safety Precaution:

1. Avoid inhalation and skin contact while diluting because of spillage or splashes.
2. Do not mix with bare hands.
3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides.
4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources.
5. The user should use full protective clothings which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat.
6. Do not smoke, eat or drink during handling and application.

Cautionary Statement:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Disposal of used containers, washing from equipment and surplus material :

It shall be the duty of manufactures, formulators of insecticides and operators to dispose packages or surplus material and washings in a safe manner so as to prevent environmental or water pollution.

- 2) The used packages shall not be left outside to prevent their re-use.
- 3) The packages shall be broken and buried away from habitation.

Manufacturer Premises :

530/2-B, Vanagaram Road, Ambattur, Chennai-600058,
Tamilnadu, India
Tamil Nadu

80/3/3, Arya Nagar & 34-36, Rural Industrial Area, Loni, District-
Ghaziabad, Uttar Pradesh, India.
Tamil Nadu

Plot No. A/74-1 & A/74-2, U.P.S.I.D.C, Sikandrabad,
Bulandshahar, Uttar Pradesh, India.
Tamil Nadu

Manufactured by

TROPICAL AGROSYSTEMS (INDIA) PVT. LTD.
72, MARSHALL'S ROAD, EGMORE CHENNAI-600008 CHENNAI
Tamil Nadu 600008

Customer Care Details:

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet