

o—|² | . . . | T . . . | qμ'® f# ! w-lw'r ! r ! # . žT' . . . -'wâ° , -' ! n p' # ° . . . Y\TWI

fi @, 3' . . . « - , -' fl

2 10 gm.mi

o—|² | . . . | T . . . | qμ'® w—| . . . « - , -' » | ° — | 3 TM qE«' fi—μ' œ—' fl—| É . . . TMμ—° 2' . . . — | ± . . . μ' . . . — a 2' 2' — 3' | ©' » Q
° 3 — | ° (œ—')

° w 3' % X 10 ⁸ </sup> " ' o³ - ©²

—° â° . . . 2^a oμ'° μμ i 0.5% w/w
° TM o—©' ® r | ' 98.50% w/w
o—|² | . . . | T . . . | qμ'® f# ! w-lw'r ! r ! # . žT' . . . -'wâ° , -' ! n p' # ° . . . Y\TWI 01.00% wp

3 TM - TSSISS' ± ' ± ' .

° μ'	— # — — 2	©' » â p ³ ' 2		® — — 2	° μ' m' a	® — — μ	® ° μ	n .
		° ' ©ñ	° ' œ—'					
qE«' fi—μ' œ—' fl	É . . . TMμ—' ff — 2 — — μ— fl	W ! " — °	o— ² . . . žT' . . . øy ³ ® — n . 3 — 2' . . . — ° a' . . . — n . 3 — 2' . . . 2 μ w ' w—° 2 — — # TM ž Y 3' — ° — ' . . . — ° — œ® — Q	-	° . p' ° ' ® μ Q	n . 3 — ©' — — » ' » Q	n . 3 — ©' — — » ' » Q	w—' . . . i ' ® œ' . . . — μ w ° . p' ° ' . . . ® μ Q

Note:- T' — μ 2' ° (œ—' 2' o—|² | . . . | — yœ' — T ! # ' 2' » Q

q[®] TM — μ w — « . . .

° | ' ® œ' — | — n . 3 — 2' . . . ©^a . . . o—|² | . . . | T . . . ž μ — , μ' ® q | ' — n . 3 — 2' . . . — w—° a' 2 μ 3 ! ° | ' ® w—μ' —
ž q 0 » Y 3' 2' . . . — 3' ©^a . . . p' — !

2 ° 2' — | a ©

2 ° 2' — | m—μ' ®' a ©³ — | — « ' — q[®] TM — 3' | — | œ » w' ©^a . . . ©—μ' « , { ' — | « ' — | o ©² μ' — | — Q

2 # | ' ® —

» μ — ° ± ' 2 # # ' © — | — μ v Q ' o ©² μ' — μ w ° Ó' ® — | — q[®] TM — Q

° μ' m' a

° . p' ° ' ® μ Q

n . « — | © —

° | ' ® œ' Q

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

° j q@œ 2 3 » a 2 3 » « © - ® - - Q

μ™ -> Q

4 gm / kg. Seed

o-|2 | . ' | ž T' | |ø³ @ - n . 3 - 2 ' - o a ' o j ' - n . 3 - 2 ' 2 μ w | ' w - o 2 - - - π™ ž Ÿ 3 ' - o - | ' - o -

° - œ - Q

n . 3 - © - » Q

- p ' o ' ® μ - m ' μ

n . 3 - © - » Q

o-|2 | . ' | ž T' | |ø³ @ - n . 3 - 2 ' - o a ' o j ' - n . 3 - 2 ' 2 μ w | ' w - o 2 - - - π™ ž Ÿ 3 ' - o - | ' - o -

° - œ - Q

n . « - - ' o 2 3

q@œ ' © ' j ' - o . p ' 2 3 ' o » - o 2 3 ' - - œ » w

w - o ' o j ' @œ - μ w . p ' o ' ® μ Q

μ™ -> » a³ - o ' qñ « - 3 μ™ j ' @œ ' 2 ' . p ' o ' ® μ - 3 j ' © - » Q

° « ž ² © μ ž qý ž j μ - | ' w μ j ' » - © - » Q

® « ® n . 1 μ ©

° © - - TMB ' 2 ' - m - o ' ® - . 1 á © - » Q

\$! m™ - - - - μ³ » © μ - @ Ÿ™™™ - qý - w Q % ³ « ñ œ | ' - @É « 1 © »™ w » © « 1 © ñ œ | ' - @É

- @ - | ' - o - - w Q & ³ « n - « 1 © »™ ' » © n - - o 2 o³ - 2 - | ' @ - o ' \$# ' o ' \$(' 2 - - μ w - w Q ! '™ -

© j ' » - ² μ j ³ Q

μ ¼ - q@œ - Q

° » - , ©

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Trichoderma viride 1.0%WP (TNAU TV-1, ITCC 6914)

(Contact Fungicide)

2 10 gm.mi

Trichoderma viride 1% WP is a bio-fungicide which effectively control root rot disease in Urd bean(black gram).

Chemical Composition:	Ingredient	%W/W
	CFU 2 X 10 ⁸ / per gm min.	
Carboxy methyl cellulose		0.5% w/w
Talc powder		98.5% w/w
Trichoderma viride 1.0%WP (TNAU TV-1, ITCC 6914)		01.00% wp
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI	Formulation					
Urd Bean(Black gram)	Root rot (Macrophomina phaseolina)	4 gm/kg of seeds	Mix required quantity of the seeds with the required quantity of Trichoderma Viride-1% WP and ensure uniform coating, shade dry and sow.	-	before sowing.	Not required	Not required	One time for seed treatment before sowing

Note:- One kilogram of formulation contains 10 gm. of Trichoderma viride culture.

Good Practices relating to pesticide application techniques

Direction of Use

Seed Treatment : Mix required quantity of the seeds with the required quantity of Trichoderma Viride-1% WP and ensure uniform coating,shade dry and sow.

Weather Condition(in general):

Should be used during favorable weather conditions and should not be used during extreme weather conditions.

Soil & Water(where ever applicable):

Light to heavy soil .Clean water to be used for application

Crop Stage for application:

before sowing.

Application Technique

Seed treatment.

Equipment Type:

Seed treatment drums or by hand wearing rubber gloves.

NozzleType:

Not Applicable.

Dosage:

Caution:Í Not to Be Used on Crops Other Than Specified on This Label/Leafletí "

4 gm / kg. Seed

Instruction for Mixing:

Mix required quantity of the seeds with the required quantity of Tricoderma Viride-1% WP and ensure uniform coating, shade dry and sow.

Re-entry Period

Not required

Pre-Harvesting Interval

Not required

Manner and Time of Application

Mix required quantity of the seeds with the required quantity of Tricoderma Viride-1% WP and ensure uniform coating, shade dry and sow.

Application Timing

sowing of the treated seed should be done in evening or early morning

Frequency

One time for seed treatment before sowing

Stages during which pesticides should not be applied on any crop :

Not applicable as the product is applied as seed treatment before sowing.

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

Symptoms Of Poisoning

Headache, Nausea, Vomiting, irritation & allergy may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to 15 minutes. 4. Remove the patient into fresh air.

Antidote

Treat Symptomatically.

Phytotoxicity

Not phytotoxic at recommended dose.

Storage Conditions

1. The packages containing the Bio-fungicides shall be stored in separate room or premises away from the rooms or premises used for storing other articles particularly articles of food or shall be kept in separate almirahs under lock and key depending upon the quantity and nature of the Bio-fungicides.
2. The rooms or premises meant for storing the Bio-fungicides shall be well built, dry, well lit and ventilated and of sufficient dimension to avoid contamination with vapour.

Safety Precaution:

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothings which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

Cautionary Statement:

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufacturers, formulators of insecticides and operators to dispose packages or surplus materials and washing in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

Village Mohiary, Chaudibagan, NH-6, Domjor, Howrah-711302
(WB)
West Bangal

Manufactured by

AMIT BIOTECH PVT. LTD.
Unit No.21, 3rd Floor, 21, Ballygunge Circular Road,
Kolkata-700019 (WB) Ballygunge West Bangal

Customer Care Details:

customer care no. : 91-33-40084152
email : admin@abplindia.com

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

o—|² l . . . | T . . . | qμ'® f# l' w-lw' r l' p' . žT' . . . -'wâ° , -'l' n p' p'° . . . Y\TWI

fi @, 3' . . . « - , -' fl

2 10 gm.mi

o—|² l . . . | T . . . | qμ'® w—| . . . « - , »' . . . —' 3 TM' qE«' fi—μ' œ—' fl—| É . . . TMμ—° 2' . . . —' ± . . . μ' . . . —' a 2' 2' —' 3' . . . | ©' . . . »Q
 ° 3' . . . —' ° (œ—' . . .

° w 3' . . . % X 10 ⁸ </sup> " . . . o³ —' ©²

—' ° â° . . . 2^a oμ'° μμ' . . . 0.5% w/w

° TM o—' ©' ® r l' . . . 98.50% w/w

o—|² l . . . | T . . . | qμ'® f# l' w-lw' r l' p' . . . 01.00% wp
 . žT' . . . -'wâ° , -'l' n p' p'° . . . Y\TWI

3 TM' . . . TSSISS' . . . ±' ±'

° μ'	—' —' —' 2	©' »â p ³ . . . 2		® —' —' 2	° μ' m' . . . a	® —' —' μ	® l' ° μ	n . . .
		° l' ©ñ	° 'œ—'					
qE«' fi—μ' œ—' fl	É . . . TMμ—' ff . . . 2 —' . . . —' j μ—' fl	W' !" —' . . . ° j	o— ² l . . . žT' . . . øy ³ ® —' n . . . 3 —' . . . 2' . . . —' ° a . . . j . . . —' . . . n . . . 3 —' 2' . . . 2 μ w . . . w—° 2 —' —' p TM ž Y 3' . . . —' ° —' . . . —' . . . ° —' œ® —' Q	-	° . p'° . . . ®μQ	n . . . 3 —' ©' —' —' »' . . . »Q	n . . . 3 —' ©' —' —' »' . . . »Q	w—' . . . —' j . . . ®œ' . . . —' μw° . . . p'° . . . ®μQ

Note:- T' —' μ 2' . . . ° œ—' 2' . . . o—|² l . . . | —' yœ' —' T!# . . . 2' . . . »Q

q^{®3} TM' —' μw —' « . . .

° j . . . ®œ' —' ° j . . . —' n . . . 3 —' 2' . . . ©^a . . . o—|² l . . . | T . . . | šμ—, μ'® q l' —' n . . . 3 —' 2' . . . —' w—° a . . . 2 μ 3!° . . . j . . . ®' w—' μ® —' . . .
žqô» Y 3' . . . 2' . . . ° —' 3' . . . ©^a . . . ° . . . p' —' !

2 ° 2' . . . —' . . . a ©

2 ° 2' . . . —' m—' μ'® . . . a ©³ . . . —' «' —' —' q^{®3} TM' —' 3' . . . j . . . —' œ »w' ©^a . . . ©—' μ' «, { . . . —' «' —' —' o ©² μ' —' —' Q

2 . . . p' | . . . ® —

»μ—' . . . ° ±' . . . 2 p p' ©—' —' μvQ' o ©² μ' —' μw° Ó'® —' —' q^{®3} TM' —' Q

° μ' m' . . . a

° . p'° . . . ®μQ

n . . . «' —' © —

° j . . . ®œ' Q

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

° j q@œ 2 3 » a 2 3 » « © - ® - - Q

μ™ -> Q

4 gm / kg. Seed

o-|2 | . ' | ž T' | |ø³ @ - n . 3 - 2 ' - o a ' o j ' - n . 3 - 2 ' 2 μ w | ' w - o 2 - - - π™ ž Ÿ 3 ' - o - | ' - o -

° - œ - Q

n . 3 - © - » Q

- p ' o ' ® μ - m ' μ

n . 3 - © - » Q

o-|2 | . ' | ž T' | |ø³ @ - n . 3 - 2 ' - o a ' o j ' - n . 3 - 2 ' 2 μ w | ' w - o 2 - - - π™ ž Ÿ 3 ' - o - | ' - o -

° - œ - Q

n . « - - ' o 2 3

q@œ ' © ' j ' - o . p ' 2 3 ' o » - o 2 3 ' - - œ » w

w - o ' o j ' @œ - μ w . p ' o ' ® μ Q

μ™ -> » a³ - o ' qñ « - 3 μ™ j ' @œ ' 2 ' . p ' o ' ® μ - 3 j ' © - » Q

° « ž ² © μ ž qý ž j μ - | ' w μ j ' » - © - » Q

® « ® n . 1 μ ©

° © - TMB ' 2 ' - m - o ' ® - . 1 á © - » Q

\$! m™ - - - - μ³ » © μ - @ Ÿ™™™ - qý - w Q % ³ « ñ œ | ' - @É « 1 © »™ w » © « 1 © ñ œ | ' - @É

- @ - | ' - o - - w Q & ' 3 « n - « 1 © »™ ' » © n - - o 2 o³ - 2 - | ' @ - o ' \$# ' o ' \$(' 2 - - μ w - w Q ! '™ -

© j ' » - 2 ' μ j ' 3 Q

μ ¼ - q@œ - Q

° » - ©

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Trichoderma viride 1.0%WP (TNAU TV-1, ITCC 6914)

(Contact Fungicide)

2 10 gm.mi

Trichoderma viride 1% WP is a bio-fungicide which effectively control root rot disease in Urd bean(black gram).

Chemical Composition:	Ingredient	%W/W
	CFU 2 X 10 ⁸ / per gm min.	
Carboxy methyl cellulose		0.5% w/w
Talc powder		98.5% w/w
Trichoderma viride 1.0%WP (TNAU TV-1, ITCC 6914)		01.00% wp
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI	Formulation					
Urd Bean(Black gram)	Root rot (Macrophomina phaseolina)	4 gm/kg of seeds	Mix required quantity of the seeds with the required quantity of Trichoderma Viride-1% WP and ensure uniform coating, shade dry and sow.	-	before sowing.	Not required	Not required	One time for seed treatment before sowing

Note:- One kilogram of formulation contains 10 gm. of Trichoderma viride culture.

Good Practices relating to pesticide application techniques

Direction of Use

Seed Treatment : Mix required quantity of the seeds with the required quantity of Trichoderma Viride-1% WP and ensure uniform coating,shade dry and sow.

Weather Condition(in general):

Should be used during favorable weather conditions and should not be used during extreme weather conditions.

Soil & Water(whenever applicable):

Light to heavy soil .Clean water to be used for application

Crop Stage for application:

before sowing.

Application Technique

Seed treatment.

Equipment Type:

Seed treatment drums or by hand wearing rubber gloves.

NozzleType:

Not Applicable.

Dosage:

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

4 gm / kg. Seed

Instruction for Mixing:

Mix required quantity of the seeds with the required quantity of Tricoderma Viride-1% WP and ensure uniform coating, shade dry and sow.

Re-entry Period

Not required

Pre-Harvesting Interval

Not required

Manner and Time of Application

Mix required quantity of the seeds with the required quantity of Tricoderma Viride-1% WP and ensure uniform coating, shade dry and sow.

Application Timing

sowing of the treated seed should be done in evening or early morning

Frequency

One time for seed treatment before sowing

Stages during which pesticides should not be applied on any crop :

Not applicable as the product is applied as seed treatment before sowing.

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

Symptoms Of Poisoning

Headache, Nausea, Vomiting, irritation & allergy may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to 15 minutes. 4. Remove the patient into fresh air.

Antidote

Treat Symptomatically.

Phytotoxicity

Not phytotoxic at recommended dose.

Storage Conditions

1. The packages containing the Bio-fungicides shall be stored in separate room or premises away from the rooms or premises used for storing other articles particularly articles of food or shall be kept in separate almirahs under lock and key depending upon the quantity and nature of the Bio-fungicides.
2. The rooms or premises meant for storing the Bio-fungicides shall be well built, dry, well lit and ventilated and of sufficient dimension to avoid contamination with vapour.

Safety Precaution:

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothings which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

Cautionary Statement:

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufacturers, formulators of insecticides and operators to dispose packages or surplus materials and washing in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

Village Mohiary, Chaudibagan, NH-6, Domjor, Howrah-711302
(WB)
West Bangal

Manufactured by

AMIT BIOTECH PVT. LTD.
Unit No.21, 3rd Floor, 21, Ballygunge Circular Road,
Kolkata-700019 (WB) Ballygunge West Bangal

Customer Care Details:

customer care no. : 91-33-40084152
email : admin@abplindia.com

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet