

Primary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10⁶/gm.min.

BRAND NAME: -

Chemical Composition :		Content % w/w
Talc powder	Talc powder	98.50% w/w
total	Total	100.00% w/w
	Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10 ⁶ /gm.min.	01.00% wp
Carboxy methyl cellulose	Carboxy methyl cellulose	0.5% w/w
Total:		100.00%

Please refer leaflet for details

Antidote

Treat Symptomatically.

Manufactured by

MSKan Biosys Pvt. Ltd.
917/17, Ravideep, Gansehwadi, Prin.K.R. Kanitkar Path, Off. F.C. Road, Pune-411004. (Maharashtra) Pune Maharashtra

Premises Address

(1). S.No. -143, Lagadmala Pune-Sinhadgad Road, Dhayri, Pune-411041 (Maharashtra) Maharashtra

Customer Care Details

sandeepakanitkar@kanbiosys.com

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10⁶/gm.min.

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

Contact Fungicide

3



Keep in cool, dry place away from heat and open flame.

Regn. No.	_____
Mfg. Lic. No.	_____
Batch No.	_____
Mfg Date	_____
Exp. Date	_____
Net Content / Volume	_____
Max. Retail Price	_____
Customercare	_____

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

3

3

Chemical Composition :		Content % w/w
Talc powder	Talc powder	98.50% w/w
total	Total	100.00% w/w
	Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10 ⁶ /gm.min.	01.00% wp
Carboxy methyl cellulose	Carboxy methyl cellulose	0.5% w/w
Total:		100.00%

3 TSS

1/4

3

3

(1).

3

sandeepakanitkar@kanbiosys.com

Safety Pictograms:



Secondary Package Label

[Ultra-small (1ml or gm to 50 ml or gm)] & [small (51 ml or gm to 250 ml or gm)]

Read leaflet before use:

Common Name:

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10⁶/gm.min.

BRAND NAME: -

Crops and Pest Groups: Trichoderma viride 1% WP is a bio-fungicide which effectively control Root rot disease in Urd Bean(Black Gram).

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For agriculture use

Re-entry:

Not Required.

Chemical Composition :

Content % w/w

Talc powder	Talc powder	98.50% w/w
total	Total	100.00% w/w
	Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10 ⁶ /gm.min.	01.00% wp
Carboxy methyl cellulose	Carboxy methyl cellulose	0.5% w/w

Total: 100.00%

Safety Precautions, Symptom of Poisoning and First Aid :Please refer leaflet for details.

Antidote Statement:

Treat Symptomatically.

Cautionary statement:

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10⁶/gm.min.

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

Contact Fungicide



Keep in cool, dry place away from heat and open flame.

Regn. No.	_____
Mfg. Lic. No.	_____
Batch No.	_____
Mfg Date	_____
Exp. Date	_____
Net Content / Volume	_____

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

Contact Fungicide

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

Keep in cool, dry place away from heat and open flame.

Talc powder	Talc powder	98.50% w/w
total	Total	100.00% w/w
	Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10 ⁶ /gm.min.	01.00% wp
Carboxy methyl cellulose	Carboxy methyl cellulose	0.5% w/w

Total: 100.00%

Safety Precautions, Symptom of Poisoning and First Aid :Please refer leaflet for details.

Antidote Statement:

Treat Symptomatically.

Cautionary statement:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Max. Retail Price _____

Manufactured by

MSKan Biosys Pvt. Ltd.
917/17, Ravideep, Gansehwadi, Prin.K.R. Kanitkar Path, Off. F.C.
Road, Pune-411004. (Maharashtra) Pune Pune Maharashtra

Premises Address

(1). S.No. -143, Lagadmala Pune-Sinhadgad Road, Dhayri,
Pune-411041 (Maharashtra)
Maharashtra

(1).

Customer Care Details

sandeepakanitkar@kanbiosys.com

sandeepakanitkar@kanbiosys.com

Safety Pictograms:



Big Label

(> 250 mL Primary and Secondary Package Label)

Read leaflet before use:

Common Name:

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10⁶/gm.min.

BRAND NAME: -

Crops and Pest Groups: Trichoderma viride 1% WP is a bio-fungicide which effectively control Root rot disease in Urd Bean(Black Gram).

Dosage, Direction for use and Manner and time of application : Please refer leaflet for details

Purpose:

For agriculture use

Re-entry:

Not Required.

Chemical Composition :

		Content % w/w
Talc powder	Talc powder	98.50% w/w
total	Total	100.00% w/w
	Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10 ⁶ /gm.min.	01.00% wp
Carboxy methyl cellulose	Carboxy methyl cellulose	0.5% w/w
Total:100.00%		

Safety Precautions:

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10⁶/gm.min.

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

Contact Fungicide

300g




Keep in cool, dry place away from heat and open flame.

Regn. No.	_____
Mfg. Lic. No.	_____
Batch No.	_____
Mfg Date	_____
Exp. Date	_____
Net Content / Volume	_____
Max. Retail Price	_____

300g

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10⁶/gm.min.

300g



300g

300g

		Content % w/w
Talc powder	Talc powder	98.50% w/w
total	Total	100.00% w/w
	Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914)CFU 2X10 ⁶ /gm.min.	01.00% wp
Carboxy methyl cellulose	Carboxy methyl cellulose	0.5% w/w
Total:100.00%		

Safety Precautions:

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothings which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

Symptom of Poisoning:

Headache, Nausea, Vomiting, irritation & allergy may occur.

First Aid:

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to 15 minutes. 4. Remove the patient into fresh air.

Antidote

Treat Symptomatically.

Cautionary statements:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Manufactured by

MSKan Biosys Pvt. Ltd.
917/17, Ravideep, Ganeshwadi, Prin.K.R. Kanitkar Path, Off. F.C. Road, Pune-411004. (Maharashtra) Pune Maharashtra

Premises Address

(1). S.No. -143, Lagadmala Pune-Sinhadgad Road, Dhayri, Pune-411041 (Maharashtra) Maharashtra

Customer Care Details

© 2017 MSKan Biosys Pvt. Ltd. All rights reserved. This document is the property of MSKan Biosys Pvt. Ltd. and is confidential. It is to be used only for the purpose for which it is intended. No part of this document may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or by any information storage and retrieval system, without the prior written permission of MSKan Biosys Pvt. Ltd.

© 2017 MSKan Biosys Pvt. Ltd. All rights reserved.

MSKan Biosys Pvt. Ltd. is a registered company in Maharashtra. The company is engaged in the business of manufacturing and marketing of various agro-chemicals. The company has a rich experience in the field of agro-chemicals and has a strong R&D facility. The company is committed to providing high quality products and services to its customers.

© 2017 MSKan Biosys Pvt. Ltd. All rights reserved.

MSKan Biosys Pvt. Ltd. is a registered company in Maharashtra. The company is engaged in the business of manufacturing and marketing of various agro-chemicals. The company has a rich experience in the field of agro-chemicals and has a strong R&D facility. The company is committed to providing high quality products and services to its customers.

© 2017 MSKan Biosys Pvt. Ltd. All rights reserved.

© 2017 MSKan Biosys Pvt. Ltd. All rights reserved.

(1).

© 2017 MSKan Biosys Pvt. Ltd. All rights reserved.

sandeepakanitkar@kanbiosys.com

Safety Pictograms:



o-|² l· |' žT' šμ-, μ'@q|' (TNAU Strain Accession no. ITCC 6914)CFU 2X10^6/gm.min.

fi @, 3' ' « - , - fl

2 x 10 gm.min

o-|² l· |' žT' |ø³' @' w- |' ' « - , »' i° - 3 TM qĒ« fi-μœ- fl- i Ē' . TMμ-° 2' - - ± .° μ' - a 2' 2' - 3' i' ©' »Q

° 3 - - ° 'œ- ' -

o-|² l· |' žT' šμ-, μ'@q|' (TNAU 01.00% wp
Strain Accession no. ITCC 6914)CFU
2X10^6/gm.min.

—° ā° ' 2 a oμ'° μμ i	—° ā° ' 2 a oμ'° μμ i	0.5% w/w
° TMō-@' @r '	° TMō-@' @r '	98.50% w/w
-μ	-μ	100.00% w/w

3 TM - TSSISS' ± '± ' -

° μ'	— x' - - 2	©' » â s³ ' 2		® - ' - 2	° μ' m' a	® - ' - μ	® l' ° μ	n .
		° !' ©ñ	° 'œ- ' -					
qĒ« fi-μœ- fl	i Ē' . TMμ- ' ff - 2 - i μ- fl		W ' ' - - i	n . 3 - © - ° ' i' ©' ' - w- ° 2 - ' ® - ° μ' ® - ' - - μw	° i' q@œ' ° . p' ° ° ' ® μQ	n . 3 - © - ' - » ' »	n . 3 - © - ' - » ' »	-

Note:- T' - μ' 2' ° 'œ- ' 2' o-|² l· |' - - yœ' \$# 2' »Q

q®³ TM' - μw - « - ' -

T' - μ' ° i' 2' w 2' fi o-|² l· |' žT' šμ-, μ'@q|' fl- 2 μ³ Q° i' ©œ' ž' i' ©' w- ° 2' - - sTM' fμ@fl- ž qō» Ÿ 3' 2' ° - - ' - ° «' . p' - Qo s©²' @' ' 2' - μw, 2' - 3' «' ' ° «' ®' 2' o ©² μ' - ž i' ° ° i' ' { ' - ° Ē- ' - μw m- μ' a ©' ©' » ° - Q o-|² l· |' žT' šμ-, μ'@q|' - n . « - - 2' ° ± .° 2 ©' ¼' 3' - - āμ' » r° ž° μ' » r° ž° - ° l' ° |' |' ' μ' ¼' 2' ° 2' - »Q

2 ° 2' - - a ©

2 ° 2' - - m- μ' ®' a ©³ ' - «' - - q®³ TM' - 3' i' - 'œ' » w' ©' - ©- μ' «, { ' - «' - - o ©² μ' - - Q

2 x |' ' © -

» μ- ° ± i' 2 s s' © - - μwQ° Ó' ® - - - o ©² μ' - Q

° μ' m' a

° i' q@œ' ° . p' ° ' ® μQ

n . « - ' © - -

° i' @œ'

° i' q@œ' ° ū° 3' ' ° - - « © -

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

n . 3 —©'—»' »

Mix 1 kg seeds with the 4 gm of Trichoderma viride 1% WP and ensure uniform coating, shade dry and sow.

—@3' "q@3 TM' — «, " «

n . 3 —©'—»' »

—p' °' @»μ' —' m©' μ

n . 3 —©'—»' »

T' —μ' °' j' 2' W' 2' fi o—|2' I' . 'j' ž' T' š' μ—, μ' @q|' fl—' 2' μ' 3' Q°' j' @œ' ž' °' j' @' w—' 2' —' m' fμ' fl—' ž' qó' Y' 3' 2' °' —' —' °' «' °' . p' —' Qo' p@2' ' @' 2' —' μw, 2' —' 3' «' °' °' «' @»' 2' o' @2' μ' —' ž' j' °' °' °' j' { —' °' É—' —' μw m—' μ' °' a' ©' @«' »' °' —' Q' o—|2' I' . 'j' ž' T' š' μ—, μ' @q|' —' n' . «' —' —' 2' °' ±' °' 2' ©' ¼' 1' 3' —' ä' μ' °' »' r' °' ž' @μ' »' r' °' ž' —' °' I' °' j' |' ' —' μ' ¼' 1' 2' °' 2' —' »' Q

n . «' —' —' °' 2' 3

°' »' 3' j' 2' —' °' 2' 3' 2' o' ©2' μ' —' 3' j' —' œ' »w

μ' TM' —' »' »' á3' —' o' °' qñ@«' —' 3' TM' °' j' @œ' °' . p' °' °' @»μ' —' 3' j' ©' »' Q

. 1' —' μ' ¼' 1'

°' «' ž' 2' ©μ' ž' qý«' ž' j' μ—' |' ' wμj' —' »' °' —' ©' »' Q

@«' ®' n . 1' μ' ©

o—|2' I' . 'j' ž' T' š' j' øy3' . @' @3' I' TM' °' °' ¼' ©' j' . —' —' , —' ž' j' °' °' °' —' TM3' 1' 2' —' m—' °' °' ®' —' . 1' á©' —' »' »' Q

n ©' ©' ©' a' —' 2' 2' μ' 2' —' j' , 3' —' —' °' @H—' Q

a) ñ œ' °' °' @H—' ž' @—' |' ' °' °' °' °' °' j' μ'

1! m' TM' —' —' , —' —' TM' μ' μ' 3' »' °' ©' TM' μ' —' @Y' TM' TM' —' qý«' —' . wQ2! 3' «' ñ œ' |' ' —' @É' «' 1' ©' »' TM' W' »' ©' «' 1' ©' ñ œ' |' ' —' @É'

—' @—' |' ' °' °' °' °' —' wQ3! 3' «' n' —' «' 1' ©' »' TM3' »' ©' n' —' —' °' 2' 63' —' 2' —' |' ' @—' °' °' 10' °' 15' 2' —' —' μw —' wQ4! TM' —'

©j' —' »' 1' 2' μ' j' 3' Q

. 1' —' —' —'

μ' ¼' 1' —' q@œ' —' —' Q

Caution:Í Not to Be Used on Crops Other Than Specified on This Label/LeafletÍ "

©

... q ... μ ... » W ... { ... « Ž w ... μ ... » ...
... Ž ... a Ž ... 3 Ž ... mō ... 2 ... ± ... μ w ... 3 ... Ž ... μ ... « ... » vQ ... 2 ... 3 ... μ ...
... ± ... » ... mčY ... » n Ž ¥ | Ž ... Ž ... « Ž ... a ... μ ... » ... w ... ± ...
... « 1 ... » ... Q ... « ... x ... » ... q ... 3 ... a ... 3 ...
... » Q

1! ... μ ... Ž ... μ ... » ... | ... « ... » ...
q ... « Ž ... | ... Y ... q ... mō ... Y ... μ w q ... μ ...
š ... mčY ... » ... vQ 4! ... » 2 ... a ... 3 ... » Ž ... μ ... ± Ž ... μ ... « 1 ... ± ...
» Ž ... » ... Y ... Q 5! q ... « ... ¼ n ... » q ... « ... » vQ ... 2 ... « ... » Ž ... 3 ... § μ ... μ ...
... 2 ... w - Ž ... 3 ... μ ... » Q 6! ... « ... » Ž ... 3 ... » Q

\$! μ ... μ ... « w ... m ... a ... mō ... » ± ... q ... μ w ... » ... » vQ ... μ ... μ ... « ... »
... 2 ... q ... « ... » ... Q

- 1! 3 ... » ... q ... μ ... » ... | ... Ž ... » ... » ... μ ... » p ... » ... ¼ ... S ...
... 1 ... » Q
- 2! q ... » w ... μ ... » ... » ... q ... » ... 3 ... » Q
- 3! q ... » w ... μ ... » ... » ... » ... » ... 2 ... » ... » Q

qñ « —

qñ « —

» « ± μ ... »

sandeepakanitkar@kanbiosys.com

¼ ... »



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10⁶/gm.min.

(Contact Fungicide)

2 x 10⁶ gm.min

Trichoderma viride 1% WP is a bio-fungicide which effectively control Root rot disease in Urad Bean(Black Gram)

Chemical Composition:	Ingredient	%W/W
	Trichoderma viride 1 % WP (TNAU Strain Accession no. ITCC 6914), CFU 2X10 ⁶ /gm.min.	01.00% wp
Carboxy methyl cellulose	Carboxy methyl cellulose	0.5% w/w
Talc powder	Talc powder	98.50% w/w
total	Total	100.00% w/w
Total:		100.00% w/w

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water	Crop Stage	Re-entry Period	Pre Harvest	Frequency
		AI	Formulation					
Urad (Black gram)	Root rot (Macrophomina Phaseolina)		4 gm/kg seed	As required for uniform coating.	Seed treatment before crop sowing	Not required	Not required	-

Note:- One kilogram of formulation contains 10 gm of Trichoderma viride biomass.

Good Practices relating to pesticide application techniques

Direction of Use

Mix 1 kg seeds with the 4 gm of Trichoderma viride (1% WP) and ensure uniform coating, shade dry and sow. Use in the evening or late afternoon to create conditions favourable for germination of spores of Trichoderma viride. The rate of application of Trichoderma viride 1 % WP is same at all confined area i.e. Glass house, Poly house, Nursery bed and Open field.

Weather Condition(in general):

Should be used during favorable weather conditions and should not be used during extreme weather conditions.

Soil & Water(where ever applicable):

Light to heavy soil .Clean water should be used for application.

Crop Stage for application:

Seed treatment before crop sowing

Application Technique

Seed Treatment

Equipment Type:

Seed treatment drums or by hand wearing rubber gloves.

NozzleType:

Not Required.

Dosage:

Mix 1 kg seeds with the 4 gm of Trichoderma viride 1% WP and ensure uniform coating, shade dry and sow.

Instruction for Mixing:

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

Please refer " Direction of use"

Re-entry Period

Not required

Pre-Harvesting Interval

Not required

Manner and Time of Application

Mix 1 kg seeds with the 4 gm of Trichoderma viride (1% WP) and ensure uniform coating, shade dry and sow. Use in the evening or late afternoon to create conditions favourable for germination of spores of Trichoderma viride. The rate of application of Trichoderma viride 1 % WP is same at all confined area i.e. Glass house, Poly house, Nursery bed and Open field.

Application Timing

Should be used preferably in early morning or evening hours

Frequency

-

Stages during which pesticides should not be applied on any crop :

Not applicable as the product is applied as seed treatment

Precautions to be taken to avoid harm to the beneficial insects like honey bees and natural enemies of pests :

Symptoms Of Poisoning

Headache, Nausea, Vomiting, irritation & allergy may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. 2. If skin and clothes are contaminated, wash the contaminated skin and clothes with plenty of water and soap. 3. If eyes are contaminated, flush the contaminated eyes with normal saline and water for 10 to 15 minutes. 4. Remove the patient into fresh air.

Antidote

Treat symptomatically

Phytotoxicity

Trichoderma viride 1% WP is eco safe bio- pesticide and Not Phytotoxicity at recommended dosage.

Storage Conditions

Store the bio-pesticide in its original pack tightly closed away from children and livestock, in a separate room, away from the rooms used for storing foodstuffs, animal feeds and other articles, under lock and key. The rooms or premises meant for storing should be well built cool dry, well ventilated, well-lit and of sufficient dimensions to avoid contamination with. Transport the bio-pesticide in such a manner so as not to come in contact with foodstuffs and animal feed

Safety Precaution:

1. Avoid inhalation and skin contact while diluting because of spillage or splashes. 2. Do not mix with bare hands. 3. After each operation, the mixing and spraying equipment is to be rinsed thoroughly with plenty of water and detergent or soap solution before being put to use for spraying other pesticides. 4. Any surplus spray solution must not be disposed off in crop lands in stagnant or flowing water, where there is a possibility of polluting water sources. 5. The user should use full protective clothings which include rubber gloves boots, either face shield or dust mask and overall rubber apron, hood or hat. 6. Do not smoke, eat or drink during handling and application.

Cautionary Statement:

1. Not to be used for any purpose other than those subscribed on label and leaflet 2. Destroy the container after use as directed on leaflet.

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "

Disposal of used containers, washing from equipment and surplus material :

1. It shall be the duty of manufacturers, formulators of insecticides and operators to dispose packages or surplus materials and washing in a safe manner so as to prevent environmental or water pollution.
2. The used packages shall not be left outside to prevent their re-use.
3. The packages shall be broken and buried away from habitation.

Manufacturer Premises :

S.No. -143, Lagadmala Pune-Sinhadgad Road, Dhayri,
Pune-411041 (Maharashtra)
Maharashtra

Manufactured by

MSKan Biosys Pvt. Ltd.
917/17, Ravideep, Gansehwadi, Prin.K.R. Kankar Path, Off. F.C.
Road, Pune-411004. (Maharashtra) Pune Maharashtra

Customer Care Details:

sandeepakanitkar@kanbiosys.com

Safety Pictograms:



Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet "