

भारत सरकार / Government of India

कृषि एवं किसान कल्याण मंत्रालय / Ministry of Agriculture & Farmers Welfare

कृषि, सहकारिता एवं किसान कल्याण विभाग / Department of Agriculture, Cooperation & Farmers Welfare

वनस्पति संरक्षण, संगरोध एवं संग्रह निदेशालय / Directorate of Plant Protection, Quarantine & Storage

एन एच 4, फरीदाबाद (हरियाणा) – 121 001/ NH-IV, Faridabad (Haryana) – 121 001

Phytosanitary condition for export of plant & plant products from India to other countries

(This compilation is NOT meant to substitute official notifications issued from time to time by the importing countries. It has been prepared ONLY for the purpose of convenient reference for general public. While efforts are made to incorporate changes from time to time, no claims/ liabilities shall be entertained for any errors that might have crept in this compilation. For authentication, relevant notification of importing countries issued may be referred to.)

Sl No.	Name of Commodity	Name of Country	Additional declaration & treatment conditions
1.	<i>Adiantum incisum</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
2.	<i>Agapanthus africanus</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
3.	<i>Agapanthus orientalis</i> Bella(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
4.	<i>Agapanthus orientalis</i> Blitza(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

5.	<i>Agapanthus orientalis</i> Maxie(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
6.	<i>Agapanthus orientalis</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
7.	<i>Agapanthus praecox</i> Black Magic (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
8.	<i>Agapanthus praecox</i> Blue Magic (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
9.	<i>Agapanthus praecox</i> Celebration(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
10.	<i>Agapanthus praecox</i> Navy Blue(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
11.	<i>Agapanthus praecox</i> Purple Cloud(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

12.	Agapanthus praecox Purple Cloud(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
13.	Agapanthus praecox Purple Delight(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
14.	Agapanthus praecox Purple Delight(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
15.	Agapanthus praecox Purple Fountain(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
16.	Agapanthus praecox Purple Magic(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
17.	Agapanthus praecox River Garden White	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
18.	Agapanthus praecox Silver Baby(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

19.	Agapanthus praecox Snowball(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
20.	Agapanthus praecox(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
21.	Agapanthus spp (Tissue Culture)	Germany	Certified that the consignment complies with Annex IV.A.I., point 32.1(d), 32.3 (d), 36.1(d), 45.1 (d) and point 46 (b) option (d) of EC Plant Health Directive 2000/29/EC
22.	Agapanthus spp(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
23.	Agapanthus spp(Tissue Culture)	United Kingdom	Certified that the consignment complies with Annex IV.A.I., point 32.1(d), 32.3 (d), 36.1(d), 45.1 (d) and point 46 (b) option (d) of EC Plant Health Directive 2000/29/EC
24.	Agave attenuata(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
25.	Agave desmetiana Variegata(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
26.	Agave desmetiana Variegata(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (Globodera rostochiensis) and (Globodera pallida) ”
27.	Agave gypsophila Blue Curls(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

28.	Agave pygmaea Dragon Toes(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
29.	Aglaonema marantifolium(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
30.	Aglaonema modestum(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
31.	Ajwain <i>Trachyspermum ammi</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
32.	Ajwain <i>Trachyspermum ammi</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
33.	Ajwain <i>Trachyspermum ammi</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
34.	Alacantarea odorata Silver(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
35.	All Fruits and Vegetables	UAE	The consignment is free from fall armyworm (<i>Spodoptera frugiperda</i>) on the basis of visual inspection
36.	Allium (Allium sp.) - Seed	Japan	This is to further certify that the parent plants were inspected during the active growth and found to be free from <i>Leptosphaeria maculans</i> (Black leg), <i>Pseudomonas viridiflava</i> (bacterial leaf blight of tomato), <i>Pseudomonas syringae</i> pv. <i>maculicola</i> (bacterial leaf spot), <i>Xanthomonas campestris</i> pv. <i>campestris</i> (black rot)
37.	Alocasia amazonica (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

38.	Alocasia amazonica Leopard (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
39.	Alocasia amazonica Variegata(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
40.	Alocasia amazonica Zebrina	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (Globodera rostochiensis) and (Globodera pallida) ”
41.	Alocasia amazonica Zebrina(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
42.	Alocasia binerva Silver Cascade(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
43.	Alocasia cuprea(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
44.	Alocasia lauterbachiana(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
45.	Alocasia lauterbachiana(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (Globodera rostochiensis) and (Globodera pallida) ”

46.	Alocasia macrorrhizos Variegata(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
47.	Alocasia sarawakensis(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
48.	Aloe affinis(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
49.	Aloe albida(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
50.	Aloe albiflora(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
51.	Aloe aristata(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
52.	Aloe humilis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
53.	Aloe spp (Tissue Culture)	Germany	Certified that the consignment complies with Annex IV.A.I., point 32.1(d), 32.3 (d), 36.1(d), 45.1 (d) and point 46 (b) option (d) of EC Plant Health Directive 2000/29/EC

54.	Aloe vera (Tissue Culture)	Netherlands/Holland	Certified that the consignment complies with point 32.1(d), 46 (b) (d) of Annex IV AI of Plant Health Directive (2000/29/EC)
55.	Aloe vera(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
56.	<i>Alpinia zerumbet Variegata</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
57.	<i>Alternanthera ocapus</i> (Tissue Culture)	Japan	Fulfills item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
58.	<i>Alternanthera reineckii</i> (Tissue Culture)	Japan	Fulfills item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
59.	Amla <i>Phyllanthusemblica</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
60.	Amla <i>Phyllanthusemblica</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
61.	Amla <i>Phyllanthusemblica</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
62.	<i>Ammomum cinnamomum</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
63.	<i>Anigozanthos</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
64.	<i>Anigozanthos viridis</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

65.	Annanas bracteatus(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
66.	Anthurium andraeanum(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
67.	Anubias barteri 'Coffeefolia' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
68.	Anubias barteri var. glabra (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
69.	Anubias barteri var. nana (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
70.	Anubias barteri var. nana (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
71.	Anubias barteri var. nana 'gold' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
72.	Anubias barteri var. nana 'petite' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
73.	Anubias heterophylla (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
74.	Apple, Pomegranate, Pear, pomegranate, Grapes, Mandarin	Srilanka	Fruits have been produced in the area free from different species of fruit flies of genera <i>Anastrepha</i> , <i>Ceratitis</i> , <i>Rhagoletis</i> , <i>Bactrocera jarvisi</i> , <i>B.musae</i> , <i>B.neohumeralis</i> , <i>B.papaya</i> , <i>B.philippinensis</i> , <i>B.tyroni</i> as verified by an official survey
75.	Arrowroot <i>Maranta arundinacea</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
76.	Arrowroot <i>Maranta arundinacea</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
77.	Arrowroot <i>Maranta arundinacea</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
78.	Aspidistra elatior(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

79.	Assorted live Plants	UAE	The consignment is free from fall armyworm (<i>Spodoptera frugiperda</i>) on the basis of visual inspection Licence No. _____ Dated: _____
80.	<i>Bacopa caroliniana</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
81.	<i>Bacopa monnieri</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
82.	<i>Bajra Pennisetum glaucum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
83.	<i>Bajra Pennisetum glaucum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
84.	<i>Bajra Pennisetum glaucum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
85.	Banana (<i>Musa</i> spp.)- Tissue culture plants	Angola	Import Permit: Certified that plantlets were raised through sterile tissue culture procedure and free of virus including Banana Bunchy Top Virus, Cucumber Mosaic virus, Banana Mosaic Virus, banana Bract Mosaic Virus, Xanthomonas (BananaWilt) using ELIZA/PCR test, Erwinia Chrysathemi, Pseudomonas Syringae Pv. Syringae (Bacterial Canker)
86.	Banana (<i>Musa</i> spp.)- Fruit	Iran	The consignment is free <i>Fromaleurodicus disperus</i> , <i>Parasaissetia nigra</i> , <i>Pentalonia nigronervosa</i> , <i>Spodoptera litura</i> , <i>Bactrocera dorsalis</i> , <i>Eudocima fullonia</i> , <i>Dysmicoccus brevipes</i> , <i>Colletrotrichum musae</i> , <i>Guignardia musae</i> , <i>Ralstonia solanacearum</i> race1,2.
87.	Basil (<i>Osamum basilicum</i>)- Seed	Kenya	The seeds was harvested from plants that have been inspected during active growth and found to be free from cucumber mosaic virus, tomato spot virus, acidovorax citrull, alfalfa mosaic virus, squash mosaic virus, zucchini yellow mosaic virus, chenephora cucurbitarum, pseudomonas syringe pv lachrymans, squash mosaic virus
88.	Basil <i>Ocimum basilicum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
89.	Basil <i>Ocimum basilicum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
90.	Basil <i>Ocimum basilicum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
91.	<i>Bauhinia purpurea</i> (Tissue Culture)	UAE	"The consignment is free from Fall Army Worm (<i>Spodoptera frugiperda</i>) on the basis of visual inspection".
92.	Bay leaf <i>Laurus nobilis</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
93.	Bay leaf <i>Laurus nobilis</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
94.	Bay leaf <i>Laurus nobilis</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
95.	<i>Begonia phoeniogramma</i> 'Type 1' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

96.	Begonia phoeniogramma 'Type 2' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
97.	Begonia roseopunctata (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
98.	Begonia roseopunctata (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
99.	Begonia spp (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
100.	Bitter gourd (<i>Momordica charantia</i>)-Seed	Bangladesh	Seeds are free from Soil and extraneous materials, and should be free from Quarantine weed seeds.
101.	Bitter gourd (<i>Momordica charantia</i>)-Seed	European Union Countries (Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech, Denmark, Estonia, Finland, Fance, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherland, Poland, Portgal, Romania, Slovakia, Slovenia, Spain, Sweden, UK)	Consignment was inspected and found free from <i>Thripspalmi</i> Karny.“Fulfils item 36.2, option two of Annex IV.A.I of Plant Health Directive (2000/29/EC) as amended.”
102.	Bitter gourd (<i>Momordica charantia</i>)-Seed	Japan	The parent plants are inspected during a field inspection and plants found free from: Acidovorax avenae pv. Citrulli, Colletotrichum orbiculare, Didymella bryoniae, Fusarium oxysporum, Parthenium hysterophorus, Pseudomonas syringae pv.Lachrymans, Squash mosaic virus, The seeds are free of trogoderma spp., Xanthomonas campestris pv.Cucurbitae, Pseudomonas pseudocaligines subsp. Citrulli, Cucumber green mottle mosaic virus. This is, further to certify that the parent plants were found free from Xanthomonas campestris pv. Cucurbitae during a field inspection.This is to further certify that the parent plants are grown from seeds disinfected against this pest or known to be free from this pest and the parent plants and fruits at a place of production are found to be free from Acidovorax avenae subsp.Citrulli by inspection during fruit maturity stage before harvesting and found to be free from Acidovorax avenae subsp.Citrulli."
103.	Bitter gourd (<i>Momordica charantia</i>)-Seed	Japan	Consignment of Plants and Plant Products is free from soil, seeds of weeds, Quarantine pests and other infection. Seeds are free from Chalara elegans, Cucumber green mottle mosaic virus,didymella bryoniae, podosphaera xanthii, setosphaeria rostrata.
104.	Bitter gourd (<i>Momordica charantia</i>)-Seed	Nepal	Seeds are free from Chalara elegans, Cucumber green mottle mosaic virus,didymella bryoniae, podosphaera xanthii, setosphaeria rostrata.

105.	Bitter gourd (<i>Momordica charantia</i>)-Seed	Vietnam	Nil
106.	Black gram <i>Vigna mungo</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
107.	Black gram <i>Vigna mungo</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
108.	Black gram <i>Vigna mungo</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
109.	Black pepper Whole <i>Piper nigrum</i> , Black pepper Powder <i>Piper nigrum</i> , Chilly Powder <i>Capsicum annum</i> , Coriander Powder <i>Coriandrum sativum</i> , Vanilla Whole <i>Vanilla planifolia</i> , Coconut Chips <i>Cocos nucifera</i> , Cardamom Seed <i>Elettaria cardamomum</i> , Coconut Chips (Raw) <i>Cocos nucifera</i> , Arjuna Powder <i>Terminalia Arjuna</i> , Amalaki Powder <i>Emblica officinalis</i> , Sariva Anantamool Powder <i>Hemidesmu indicus</i> , Bala Powder <i>sida cordifolia</i> , Rama Tulsi Powder FFL <i>Ocimum tenuiflorum</i> , Turmeric Powder <i>Curcuma Longa</i> , Haritaki Powder FW <i>Terminalia chebula</i> , Kapikachu Powder <i>Mucuna pruriens</i> , Punarnava Powder <i>Boerhaavia diffusa</i> , Triphala Powder (FW) <i>Multi ingredient product</i> , Vidhari Khanda <i>Ipomoea digitata</i> , Bilva Powder <i>Aegle marmelos</i> , Brahmi <i>Bacopa Monnieri</i> , Musta Powder <i>Cyperus Rotundus</i> , Guduchi Powder <i>Tinospora Cordifolia</i>	Canada	NIL

110.	Blechnum gibbum (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
111.	Bottle Gourd- (Lagenaria siceraria)- Seed	Japan	The parent plants or the seeds harvested from the parents plants are tested by an appropriate genetic method such as RT-PCR assay and found to be free from Potato spindle tuber viroid. For seed test, a sample of 4,600 seeds randomly drawn from lot in accordance with the International Seed Testing Association (ISTA) procedures is divided and tested as sub-samples of no more than 400 seeds for RT-PCR assay
112.	Braccoli (<i>Brassica oleracea var. italic</i>)- Seed	Bulgaria	Nil
113.	Braccoli (<i>Brassica oleracea var. italic</i>)- Seed	Peru	Nil
114.	Brunfelsia latifolia (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
115.	Bucephalandra kishii (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
116.	Bucephalandra sp. ‘sintang’ (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
117.	Bucephalandra sp. 'Sintang' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
118.	Byadgi Chilli Whole <i>Capsicum annum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
119.	Byadgi Chilli Whole <i>Capsicum annum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
120.	Byadgi Chilli Whole <i>Capsicum annum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
121.	Cabbage (<i>Brassica oleracea Var. capitata</i>)- Seeds-	Bangladesh	Seeds are free from Bacterial leaf Blight of tomato (<i>Pseudomonas viridiflava</i>), Bacterial Spot (<i>Pseudomonas syringae</i> pv). Block rot (<i>Xanthomonas Campestris</i> pv), Black leg (<i>Leptosphaeria maculans</i>). Seeds are free from soil and extraneous materials, and quarantine weed seed
122.	Cabbage (<i>Brassica oleracea Var. capitata</i>)- Seeds-	Bangladesh	Plants and plant products are free from injurious insect pest and disease, extraneous materials and quarantine weed seeds. Seeds are free from (a) Black leg (<i>Leptosphaeria maculans</i>) (b)Bacterial leaf blight (<i>Pseudomonas viridiflava</i>) (c)Bacterial spot (<i>Pseudomonas syringae</i> pv) (d)Black rot(<i>Xanthomonas campestris</i> pv) . Seed are treated with Thiram
123.	Cabbage (<i>Brassica oleracea Var. capitata</i>)- Seeds-	Jamaica	Sees are free from <i>Leptosphaeria maculans</i> and <i>Xanthomonas campestris</i> pv raphani.

124.	Cabbage (<i>Brassica oleracea</i> Var. <i>capitata</i>)- Seeds-	Jamaica	Sees are free from <i>Leptosphaeria maculans</i> and <i>Xanthomonas campestris</i> pv <i>raphani</i> .
125.	Cabbage (<i>Brassica oleracea</i> Var. <i>capitata</i>)- Seeds-	Kenya	The seeds were tested and found to be free from the pests <i>Rhodococcus fascians</i> (Fasciation: leafy gall) <i>Xanthomonas campestris</i> pv <i>campestris</i> , Aster yellow beet mosaic virus, cauliflower mosaic virus and turnip mosaic virus Seeds are treated with Thiram @ 2gm /kg of seed
126.	Cabbage (<i>Brassica oleracea</i> Var. <i>capitata</i>)- Seeds-	Mauritius	(<i>Brassica oleracea capitata</i>) Free from (a) <i>Xanthomonas campestris</i> pv. <i>Campestris</i> (b) <i>Leptosphaeria maculans</i> (c) <i>Pseudomonas viridiflava</i> (d) <i>Pseudomonas syringae</i> pv <i>maculicola</i>
127.	Cabbage (<i>Brassica oleracea</i> Var. <i>capitata</i>)- Seeds-	Zambia	Free from <i>albugo candida</i> , <i>botrytinia fuckelina</i> , <i>peronospora parasitica</i> , <i>gibberella avenacea</i> and <i>lolium temulentam</i> and orabanche and fungicide treatment
128.	<i>Calathea ornata</i> 'beauty star' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
129.	<i>Callicostella prabaktiana</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
130.	<i>Callicostella prabaktiana</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
131.	<i>Callistemon viminalis</i> (Tissue Culture)	Australia	"Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened". AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
132.	<i>Canna Tropicana</i> (Tissue Culture)	Australia	"Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened". AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
133.	Cantaloupe (<i>Cucumis melo</i> var. <i>cantalupensis</i>)- Seed	Jamaica	Seeds are free from <i>Septoria</i> sp.(<i>Septoria</i> Leaf Spot)
134.	<i>Cardamine lyrata</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
135.	<i>Cassia siamea</i> (Tissue Culture)	UAE	"The consignment is free from Fall Army Worm (<i>Spodoptera frugiperda</i>) on the basis of visual inspection".
136.	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)- Seed	Bangladesh	Seeds are free from diseases and insect a. Bacterial Leaf Blight of tomato (<i>Pseudomonas viridiflava</i>) b. Bacterial Spot (<i>Pseudomonas syringae</i> pv) c. Block rot (<i>Xanthomonas Campestris</i> pv) d. Black leg (<i>Leptosphaeria maculans</i>) e. Leaf miner (<i>Liriomyza trifolii</i>) Seeds are free from soil and extraneous materials, and quarantine weed seed.
137.	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)- Seed	Japan	This is to further certify that the parent plants were inspected during the active growth and found to be free from <i>Leptosphaeria maculans</i> (Black leg) , <i>Pseudomonas viridiflava</i> (bacterial leaf blight of tomato) , <i>Pseudomonas syringae</i> pv. <i>maculicola</i> (bacterial leaf spot) , <i>Xanthomonas campestris</i> pv. <i>campestris</i> (black rot)

138.	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)- Seed	Japan	This is to further certify that the parent plants were inspected during the active growth and found to be free from <i>Leptosphaeria maculans</i> (Black leg), <i>Pseudomonas viridiflava</i> (bacterial leaf blight of tomato), <i>Pseudomonas syringae</i> pv. <i>maculicola</i> (bacterial leaf spot), <i>Xanthomonas campestris</i> pv. <i>campestris</i> (black rot)
139.	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)- Seed	Japan	Consignment of Plants and Plant Products is free from soil, seeds of weeds, Quarantine pests and other infection. Seeds are free from <i>Gibberella avenacea</i> , <i>leptosphaeria maculans</i> , <i>phytophthora cryptogea</i> , <i>verticillum albo-atrum</i> , <i>erwina carotovora</i> subsp. <i>atroseptica</i> , <i>pseudomonas cichorii</i> , <i>rhodococcus fascians</i> , beet western yellow virus, broccoli necrotic yellow virus, cauliflower mosaic virus,
140.	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)- Seed	Nepal	Seeds are free from <i>Gibberella avenacea</i> , <i>leptosphaeria maculans</i> , <i>phytophthora cryptogea</i> , <i>verticillum albo-atrum</i> , <i>erwina carotovora</i> subsp. <i>atroseptica</i> , <i>pseudomonas cichorii</i> , <i>rhodococcus fascians</i> , beet western yellow virus, broccoli necrotic yellow virus, cauliflower mosaic virus,
141.	Cauliflower (<i>Brassica oleracea</i> var. <i>botrytis</i>)- Seed	South Africa	Free from Tobacco ring spot nepovirus, tomato ring spot nepovirus and squash mosaic virus
142.	Cauliflower (<i>Brassica oleracea</i>) Seeds	West Indies	Seeds are free from soil, Trash, Organic Matter, Plant Pests, and Disease. Seeds treated with a Fungicide before shipping
143.	Chilli (<i>Capsicum annum</i>)- Seed	Bangladesh	Seeds have been collected from strand free from chilli mosaic virus anthracnose (<i>Colletotrichum tematium</i>), free from following pepper viruses (i) Mild mosaic virus (ii) Mild mottle virus (iii) Tomato ring spot virus (iv) Tomato black ring virus. Seeds are free from bacterial scab (<i>Xanthomonas vesicatoria</i>) and quarantine weed seeds.. Seeds treated with Polymer + Thiram Export vide Import Permit Number XXXXX Dated XXXX
144.	Chilli (<i>Capsicum annum</i>)- Seed	Japan	The parent plants or the seeds harvested from the parents plants are tested by an appropriate genetic method such as RT-PCR assay and found to be free from Potato spindle tuber viroid. For seed test, a sample of 4,600 seeds randomly drawn from lot in accordance with the International Seed Testing Association (ISTA) procedures is divided and tested as sub-samples of no more than 400 seeds for RT-PCR assay
145.	Chilli (<i>Capsicum annum</i>)- Seed	Tanzania	Nil
146.	Chilli (<i>Capsicum annum</i>)- Seed	West Indies	Seeds are free from soil, Trash, Organic Matter, Plant Pests, and Disease. Seeds treated with a Fungicide before shipping
147.	Chilli (<i>Capsicum annum</i>)- seed	Bulgaria	Nil
148.	Chilli (<i>Capsicum frutescens</i>)- Seed	Bangladesh	The consignment is free from injurious pests, & the seeds have been collected from strand free from chilli mosaic virus anthracnose (<i>Colletotrichum tema um</i>) & for freedom from following Pepper viruses, Mild mosaic virus., Mild Motle virus., Tomato ring spot virus. Tomato blacking virus. The Consignment should be free from bacterial Scab (<i>Xanthomonas Vesicatoria</i>), The Consignments shall be free from quarantine weed seeds.
149.	Chilli, Sweet & Hot pepper (<i>Capsicum annum</i>) Seed	Iran	The seeds were free from of pest and disease mentioned below : <i>Colletotricum capsii</i> , <i>ralstonia solanacearum</i> , <i>xanthomonas vesicatoria</i> Seeds were treated with fungicide and free from soil, plant material, live insects, and plant diseases and weed seeds
150.	Chilli/ Pepper (<i>Capsicum frutescens</i>)- Seed	South Korea	Nil

151.	Chilli/ Sweet pepper/ Capsicum (<i>Capsicum frutescens</i>)- Seed	USA	Seeds were found free from <i>Clavibacter michiganensis</i> ssp. <i>Michiganensis</i> , <i>Xanthomonas</i> (<i>campestris</i> pv.) <i>vesicatoria</i> , Potato spindle tuber viroid, Pepino mosaic virus
152.	Chilli/Bell Pepper (<i>Capsicum annuum</i>)- Fresh Fruit	UAE	The lot of Bell Pepper accepted according to the codex alimentarius maximum residue limits vide residue analysis report number #VLL/BLR/18-19/1593 14.08.2018 issued by Vimta labs ltd., RR District, Hyderabad (quality@vimta.com), India as per requirement of ministry of agriculture of UAE as certificate attached in original
153.	Chives (<i>Allium schoenoprasum</i>)-Fresh Vegetables	European Union	NIL
154.	<i>Chlorophytum comosum</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
155.	<i>Clivia miniata</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
156.	Cloves <i>Syzygium aromaticum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
157.	Cloves <i>Syzygium aromaticum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
158.	Cloves <i>Syzygium aromaticum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
159.	<i>Colocasia antiquarum</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
160.	<i>Colocasia heterochroma</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

161.	Cordyline australis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
162.	Cordyline australis Black Knight (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
163.	Cordyline australis Burgandy (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
164.	Cordyline australis Green Magic (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
165.	Cordyline australis Kermit (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
166.	Cordyline australis New Red(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
167.	Cordyline australis Pink Variegated (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

168.	Cordyline australis Pocahontas(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
169.	Cordyline australis Radiant (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
170.	Cordyline australis Red Sensation(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
171.	Cordyline australis Red Sensation(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
172.	Cordyline australis Red Sensation(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
173.	Cordyline australis Red Star (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
174.	Cordyline australis Red Star(Tissue Culture)	Canada	The plants were produced by <i>InVitro techniques</i> and have never been in contact with soil, soil related matter or growing media outside the laboratory.
175.	Cordyline australis Rededge (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

176.	Cordyline australis Torbay Dazzler (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
177.	Cordyline fruticosa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
178.	Cordyline Fruticosa Black	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
179.	Cordyline fruticosa Black(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
180.	Cordyline fruticosa John Klass(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
181.	Cordyline fruticosa Kiwi (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
182.	Cordyline fruticosa Kiwi (Tissue Culture)	Netherlands/Holland	Certified that the consignment complies with point 32.1(d), 46 (b) (d) of Annex IVAI of Plant Health Directive (2000/29/EC)
183.	Cordyline fruticosa New Guinea Fan (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

184.	Cordyline fruticosa Schuberti (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
185.	Cordyline fruticosa Tangelo (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
186.	Cordyline spp (Tissue Culture)	Germany	Certified that the consignment complies with Annex IV.A.I., point 32.1(d), 32.3 (d), 36.1(d), 45.1 (d) and point 46 (b) option (d) of EC Plant Health Directive 2000/29/EC
187.	Cordyline spp (Tissue Culture)	USA	Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (Globodera rostochiensis) and (Globodera pallida) ”
188.	Cordyline terminalis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
189.	Cordyline terminalis Black Magic (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
190.	Cordyline terminalis Blue Magic (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
191.	Cordyline terminalis Brayand Diamond (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

192.	Cordyline terminalis Bronze (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
193.	Cordyline terminalis Bronze(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
194.	Cordyline terminalis Chocolate Queen (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
195.	Cordyline terminalis Chocolate Queen (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
196.	Cordyline terminalis Dr.Brown(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
197.	Cordyline terminalis Early Morning Diamond(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
198.	Cordyline terminalis Green Isle(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

199.	Cordyline terminalis Hilo Rainbow(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
200.	Cordyline terminalis Mahatma(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
201.	Cordyline terminalis Maize(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
202.	Cordyline terminalis Maize(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (Globodera rostochiensis) and (Globodera pallida) ”
203.	Cordyline terminalis Miss Jules (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
204.	Cordyline terminalis Negra (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
205.	Cordyline terminalis Negra(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
206.	Cordyline terminalis Pink Chocolate (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

207.	Cordyline terminalis Pink Diamond (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
208.	Cordyline terminalis Pink Variegated (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
209.	Cordyline terminalis Red	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
210.	Cordyline terminalis Red Sister Gold Edge (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
211.	Cordyline terminalis Rededge (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
212.	Cordyline terminalis Rededge(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
213.	Cordyline terminalis Rubra (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

214.	Cordyline terminalis Ruby (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
215.	Cordyline terminalis Tropic Snow(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
216.	Cordyline terminalis Cream Delight (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
217.	Cordyline terminalis Green (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
218.	Cordyline terminalis Green Isle(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
219.	Cordyline terminalis Pink Delight (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
220.	Cordyline terminalis Purple Princess (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

221.	Cordyline terminalis Red Delight (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
222.	Cordyline terminalis Tikki (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
223.	Cordyline terminalis White Maginata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
224.	Cordyline terminalis ZIGZAG (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
225.	Coriander (Coriandrum sativum)- Seed	Kenya	The seeds was harvested from plants that have been inspected during active growth and found to be free from cucumber mosaic virus, tomato spot virus, acidovorax citrull, alfalfa mosaic virus, squash mosaic virus, zucchini yellow mosaic virus, and cucumber mosaic virus Seeds are treated with thiram 2 g and 5 ml red polymer per kg seed before dispatch
226.	Coriander (Coriandrum sativum)- Seed	Nepal	Consignment of Plants and Plant Products is free from soil, seeds of weeds and other infection.
227.	Coriander <i>Coriandrum sativum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
228.	Coriander <i>Coriandrum sativum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
229.	Coriander <i>Coriandrum sativum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
230.	Cotton, Gossypium spp.	Pakistan	The consignment of raw cotton is free from all the insect pests and diseases particularly <i>Anthonomus grandis</i> . Fumigation with methyl bromide (mbr) @ 80grams/m3 for 48 hrs
231.	Cotton, Gossypium spp.	Indonesia	The consignment of cotton is free from all the insect pests and diseases particularly <i>Anthonomus grandis</i> . Fumigation with methyl bromide (mbr) @ 80grams/m3 for 48 hrs

232.	Cotton, <i>Gossypium</i> spp.	China	The cotton is free from cotton boll weevils <i>Anthonomus grandis</i> , <i>Anthonomus peninsularis</i> and <i>Anthonomus vestitus</i> . Fumigation with Methyl Bromide (mbr) @ 80GRAMS/M3 for 48 Hrs
233.	Cotton, <i>Gossypium</i> spp.	Turkey	Cotton is free from plant material and cotton seeds.
234.	Cotton, <i>Gossypium</i> spp.	Bangladesh	This consignment is free from injurious insect pest and disease. Fumigation with Methyl Bromide (MBr) @ 48 GRAMS/M3 for 24 Hrs
235.	Cotton, <i>Gossypium</i> spp.	Vietnam	The consignment is free from khapra beetle (<i>Trogoderma granarium</i>). Fumigation with Methyl Bromide (MBr) @ 100 GRAMS/M3 for 74 Hrs
236.	<i>Crimum pedunculatum</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
237.	<i>Cryptocoryne axelrodi</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
238.	<i>Cryptocoryne crispatula</i> var. <i>balansae</i> 'Green' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
239.	<i>Cryptocoryne spiralis</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
240.	<i>Cryptocoryne spiralis</i> 'Tiger' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
241.	<i>Cryptocoryne wendtii</i> 'Green' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
242.	<i>Cryptocoryne wendtii</i> 'Tropica' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
243.	<i>Ctenanthe pilosa</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
244.	<i>Ctenanthe pilosa</i> Golden Mozaic (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

245.	Ctenanthe setosa (Tissue Culture)	Australia	"Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened". AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
246.	Ctenanthe setosa Grey Star (Tissue Culture)	Australia	"Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened". AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
247.	Cucumber	Taiwan	Seeds had been thoroughly inspected and found free from spindle tuber viroid based on official laboratory analysis
248.	Cucumber (<i>Cucumis sativus</i>)- Seed	Japan	"Fulfills item 19 of the Annexed Table 2-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No73/1950)"
249.	Cucumber (<i>Cucumis sativus</i>)- Seed	Jordan	Nil
250.	Cucumber (<i>Cucumis sativus</i>)- Seed	Republic of Korea	Nil
251.	Cucumber (<i>Cucumis sativus</i>)- Seeds	Japan	The parent plants are grown from seeds disinfected against <i>Acidovorax avenae</i> subsp. citrulli or known to be free from this pest AND The parent plants and fruits at a place of production or a production site (including a plant growth facility) are found to be free from <i>Acidovorax avenae</i> subsp.citrulli by inspection, including laboratory testing of any suspicious symptoms, carried out during fruit maturity stage before harvesting.
252.	Cucumber (<i>Cucumis sativus</i>)- Seeds	Thailand	Nil
253.	Cucumber-(<i>Cucumis sativus</i> -) Seeds	Japan	The parent plants are grown from seeds disinfected against this pest or known to be free from this pest And The parent plants and fruits at a place of production or a production site (including a plant growth facility) are found to be free from <i>Acidovorax avenae</i> subsp. citrulli by inspection, including laboratory testing of any suspicious symptoms, carried out during fruit maturity stage before harvesting. This is to further certify that the parent plants were inspected during the active growth and found to be free from Cucumber leaf spot virus, Arabis mosaic virus, <i>Didymella bryoniae</i> .
254.	Cumin <i>Cuminum cyminum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
255.	Cumin <i>Cuminum cyminum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
256.	Cumin <i>Cuminum cyminum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
257.	Cumin Powder <i>Cuminum cyminum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
258.	Cumin Powder <i>Cuminum cyminum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.

259.	Cumin Powder <i>Cuminum cyminum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
260.	<i>Curcuma alismatifolia</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
261.	<i>Curcuma alismatifolia</i> Variegata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
262.	<i>Curcuma longa</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
263.	Curry leaf dry <i>Murraya koenigii</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
264.	Curry leaf dry <i>Murraya koenigii</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
265.	Cut flowers	Turkey	Immediately prior to their export, they have been officially inspected and found free from <i>Bemisia tabaci</i> “the consignment complies with decree no. 29345 published on 04.05.2015 annex iv item 56.3
266.	Cut flowers	Japan	Immediately prior to their export, have been officially inspected and found free from <i>Circulifer tenellus</i> . “fulfills item 9 of the annexed table 2-2 of the ordinance for enforcement of the plant protection act. ((maf ordinance no 73 /1950)
267.	Cut flowers	Mauritius	Plants are free from san jose scale (<i>Quadraspidiotus perniciosus</i>) crown call (<i>Agrobacterium tumefactens</i>), rose mosaic virus, rose rosette virus , strawberry latent ring spot virus, rose wilt disease and tomato ringspot virus/plants are treated with fungicide (carbendazim 50% w.p.)and insecticides (imidacloprid 80%)
268.	Curry leaf dry <i>Murraya koenigii</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
269.	<i>Delonix regia</i> (Tissue Culture)	UAE	"The consignment is free from Fall Army Worm (<i>Spodoptera frugiperda</i>) on the basis of visual inspection".
270.	<i>Dendranthema</i> (<i>Chrysanthemum sub set Dendranthema sp.</i>) Cut Flowers	European Union	Fulfils item 32.2, option two of Annex IVAI of Plant Health Directive (2000/29/EC) as amended. Immediately prior to their export, have been officially inspected and found free from <i>Liriomyza sativae</i> (Blanchard) and <i>Amauromyza maculosa</i> (Malloch).

271.	Dhatara (<i>Datura stramonium</i>) (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
272.	Dhatara (<i>Datura stramonium</i>) (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
273.	Dhatara (<i>Datura stramonium</i>) (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
274.	Dianella (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
275.	Dianella caerulea Green(Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IHR, Bangalore-Govt.of India
276.	Dianella caerulea Lucia (Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IHR, Bangalore-Govt.of India
277.	Dianella revoluta (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
278.	Dianella revoluta Destiny (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

279.	Dianella revoluta Rainbow Twist (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
280.	Dianella revoluta Silver Streak (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
281.	Dianella revoluta Silver streak(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
282.	Dianella revoluta Variegata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
283.	Dianella revoluta Wyeena (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
284.	Dianthus (<i>Dianthus sp.</i>) Cut Flowers	European Union	Fulfils item 32.2, option two of Annex IVAI of Plant Health Directive (2000/29/EC) as amended. Immediately prior to their export, have been officially inspected and found free from Liriomyza sativae (Blanchard) and Amauromyza maculosa (Malloch).
285.	Dieffenbachia maculata Camille (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
286.	Dieffenbachia maculata Compacta (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

287.	Dieffenbachia maculata Perfecta	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
288.	Dieffenbachia maculata Sparkles (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
289.	Dieffenbachia maculata Tropic Snow (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
290.	Dieffenbachia maculata White Blizzard (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
291.	Dieffenbachia maculata(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (Globodera rostochiensis) and (Globodera pallida) ”
292.	Dieffenbachia maculate (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
293.	Dionea (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
294.	Diplazium cordifolium (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

295.	Dracaena deremensis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
296.	Dracaena massangeana (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
297.	Dracaena massangeana Highlighter (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
298.	Dracaena massangeana White Aspen (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
299.	Dracaena massangeana White Aspen(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (Globodera rostochiensis) and (Globodera pallida) ”
300.	Dracaena sanderiana (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
301.	Dry Ginger <i>Zingiber officinale</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
302.	Dry Ginger <i>Zingiber officinale</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
303.	Dry Ginger <i>Zingiber officinale</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
304.	Dry Whole Coconut <i>Coconus nucifera</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
305.	Dry Whole Coconut <i>Coconus nucifera</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.

306.	Dry Whole Coconut <i>Coconus nucifera</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
307.	Dry Whole Coconut <i>Coconus nucifera</i> (Food Product)	Iran	The consignment is free from <i>Asterole canium pustulans</i> , <i>Aleurodicus distractor</i> , <i>Dysmicoccus brevipes</i> , <i>Nipaecoccus nipae</i> , <i>Parasaissetia nigra</i> , <i>Unaspis citri</i> , <i>Rhynchophorus pal marum</i> , <i>Rhynchophorus ferrugineus</i> , <i>Araecerus fasciculatus</i> , <i>Tirathaba rufivena</i> , <i>Iceria pulchra</i> .
308.	Echevaria agavoides (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
309.	Echevaria agavoides(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (<i>Globodera rostochiensis</i>) and (<i>Globodera pallida</i>) ”
310.	Echevaria australis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
311.	Echevaria elegans	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (<i>Globodera rostochiensis</i>) and (<i>Globodera pallida</i>) ”
312.	Echevaria elegans (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
313.	Echevaria longissima (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
314.	Echevaria nodulosa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

315.	Echeveria nodulosa(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (<i>Globodera rostochiensis</i>) and (<i>Globodera pallida</i>)”
316.	Echeveria agavoides Purple Pearl (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
317.	Echeveria derenbergii(Tissue Culture)	Kenya	Certified that the live tissue culture plants in this consignment were propagated and grown in a sterile medium
318.	Echeveria desmetiana (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
319.	Echeveria elegans Blue (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
320.	Echeveria glauca (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at
321.	Echeveria globulosa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
322.	Echeveria minima (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

323.	Echeveria paniculata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
324.	Echeveria pulvinata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
325.	Echeveria pulvinata(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (<i>Globodera rostochiensis</i>) and (<i>Globodera pallida</i>) ”
326.	Echeveria racemosa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
327.	Echeveria setosa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
328.	Echeveria spp (Tissue Culture)	Germany	Certified that the consignment complies with Annex IV.A.I., point 32.1(d), 32.3 (d), 36.1(d), 45.1 (d) and point 46 (b) option (d) of EC Plant Health Directive 2000/29/EC
329.	Echeveria spp Red Edge (Tissue Culture)	Kenya	Certified that the live tissue culture plants in this consignment were propagated and grown in a sterile medium
330.	Echinodorus rubin (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
331.	Echinodorus argentinensis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
332.	Echinodorus argentinensis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
333.	Echinodorus arjuna (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
334.	Echinodorus grandiflorus (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
335.	Echinodorus grandiflorus (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

336.	Echinodorus joyo 'red star' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
337.	Echinodorus marble queen (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
338.	Echinodorus ozelot 'Green' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
339.	Echinodorus ozelot 'Green long leaf'	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
340.	Echinodorus tanzadefuder (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
341.	Echinodorus tanzadefuder (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
342.	Egg Plant (<i>Solanum melongena</i>)- Seed	Jordan	Nil
343.	Eleocharis acicularis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
344.	Eleocharis acicularis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
345.	Eleocharis parvula (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
346.	Eleocharis parvula (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
347.	Elettaria spp (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
348.	Epiprenum pinnatum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
349.	Fennel <i>Foeniculum vulgare</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
350.	Fennel <i>Foeniculum vulgare</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
351.	Fennel <i>Foeniculum vulgare</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.

352.	Fennel seeds <i>Foeniculum vulgare</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
353.	Fennel seeds <i>Foeniculum vulgare</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
354.	Fennel seeds <i>Foeniculum vulgare</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
355.	Fern (Tissue Culture)	USA	Tissue cultures in this consignment has been grown <i>InVitro</i> , and produced within a secure environment in a production area that is free of Potato cyst nematodes and has never been grown in soil nor come in contact with soil, as indicated on laboratory test dated XXXXX29/01/2019 IIHR, Bangalore-Govt.of India.
356.	Ficus benghalensis Audrey (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
357.	Ficus benghalensis Variegated (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
358.	Ficus elastica Burgandy (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
359.	Ficus elastica Ruby (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
360.	Ficus elastica Shivereana (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

361.	Ficus elastica Shivereana(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
362.	Ficus elastica Tineke (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
363.	Ficus lyrata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
364.	Ficus lyrata Bambino (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
365.	Ficus lyrata Bambino(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
366.	Flax seeds <i>Linum usitatissimum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
367.	Flax seeds <i>Linum usitatissimum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
368.	Flax seeds <i>Linum usitatissimum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
369.	Food products- Rice Grains/ Flour	USA	This shipment has been inspected and found free of khapra beetle (trogoderma granarium) Container no. XXXX Inspection conducted on 15.06.2018
370.	Fried onion <i>Allium cepa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
371.	Fried onion <i>Allium cepa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.

372.	Fried onion <i>Allium cepa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
373.	Garlic bulb <i>Allium sativum</i> (Food Product)	Taiwan	The consignment is free from <i>Rhizoglyphus echinopus</i> , <i>Ditylenchus dipsaci</i> , <i>Delia antiqua</i> , <i>Delia platura</i> , <i>Petrobia lateens</i> , <i>Stromatinia cepivora</i> (<i>Sclerotium cepivorum</i>), <i>Urocystis cepulae</i> , <i>Chenopodium murale</i> .”
374.	Gasteria armstrongii (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
375.	Gasteria batesiana (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
376.	Gasteria spp (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
377.	Gasteria spp Arack(Tissue Culture)	Kenya	Certified that the live tissue culture plants in this consignment were propagated and grown in a sterile medium
378.	Gasteria spp Arack(Tissue Culture)	Kenya	Certified that the live tissue culture plants in this consignment were propagated and grown in a sterile medium
379.	Gasteria spp Green Ice(Tissue Culture)	Kenya	Certified that the live tissue culture plants in this consignment were propagated and grown in a sterile medium
380.	Gasteria spp Thais(Tissue Culture)	Kenya	Certified that the live tissue culture plants in this consignment were propagated and grown in a sterile medium
381.	Glossostigma elatinoides	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
382.	Glossostigma elatinoides (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
383.	Goldenrods (<i>Solidago sp.</i>) Cut Flowers	European Union	Fulfils item 32.2, option two of Annex IVAI of Plant Health Directive (2000/29/EC) as amended. Immediately prior to their export, have been officially inspected and found free from <i>Liriomyza sativae</i> (Blanchard) and <i>Amauromyza maculosa</i> (Malloch).
384.	Gratiola viscidula (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
385.	Green Coriander Powder <i>Coriandrum sativum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.

386.	Green Coriander Powder <i>Coriandrum sativum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
387.	Green Coriander Powder <i>Coriandrum sativum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
388.	Grevillea longistyla (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
389.	Grevillea longistyla Carpet Layer (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
390.	Grevillea longistyla Fire Sprite (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
391.	Grevillea longistyla Pinky Petite (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
392.	Grevillea robusta (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
393.	Grevillea rosmarinifolia (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

394.	Grevillea rosmarinifolia Bronze Ramble (Tissue Culture)r	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
395.	Grevillea rosmarinifolia Coconut Ice (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
396.	Grevillea rosmarinifolia Deuo Flame (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
397.	Grevillea rosmarinifolia Forest Rambler (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
398.	Grevillea rosmarinifolia Gold Rush (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
399.	Grevillea rosmarinifolia Honey Gem (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
400.	Grevillea rosmarinifolia Jester (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

401.	Grevillea rosmarinifolia Lana Maree (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
402.	Grevillea rosmarinifolia Little Robyn (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
403.	Grevillea rosmarinifolia Loopy Lou (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
404.	Grevillea rosmarinifolia Moonlight (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
405.	Grevillea rosmarinifolia Mt Tamborintha (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
406.	Grevillea rosmarinifolia Ned Kelly (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
407.	Grevillea rosmarinifolia Robyn Gordon (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

408.	Grevillea rosmarinifolia Royal Mantle (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
409.	Grevillea rosmarinifolia Sandra Gorden (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
410.	Grevillea rosmarinifolia Soopa Doopa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
411.	Grevillea rosmarinifolia Sun Kissed (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
412.	Grevillea rosmarinifolia Superb (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
413.	Gypsophila (<i>Gypsophila sp.</i>)- Cut Flowes	European Union	Fulfils item 32.2, option two of Annex IVAI of Plant Health Directive (2000/29/EC) as amended. Immediately prior to their export, have been officially inspected and found free from Liriomyza sativae (Blanchard) and Amauromyza maculosa (Malloch).
414.	Habranthus robusta (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
415.	Haworthia attenuate (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

416.	Haworthia coarctata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
417.	Haworthia fasciata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
418.	Haworthia limifolia (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
419.	Haworthia marginata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
420.	Haworthia retusa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
421.	Haworthia venusta (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
422.	Hemianthus callitrichoides 'Cuba' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
423.	Hemianthus micranthemoides (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
424.	Hemionitis ariforia (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
425.	Homalomena humilis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

426.	Homalomena rubescens (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
427.	Hosta (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
428.	Hosta fortune (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
429.	Hydrocotyle leucocephala (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
430.	Hydrocotyle tripartite (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
431.	Hydrocotyle verticillata (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
432.	Hygrophila angustifolia (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
433.	Hygrophila difformis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
434.	Hygrophila lacustris (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
435.	Hygrophila pinnatifida (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
436.	Hygrophila pinnatifida (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
437.	Hygrophila pinnatifida (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
438.	Hygrophila polysperma (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
439.	Hygrophila sp 'Tiger' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
440.	Hygrophila stricta (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

441.	Hymenocallis littoralis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
442.	Hymenocallis littoralis Variegated (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
443.	Iris tectorum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
444.	Ixora accuminata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
445.	Kalanchoe orgyalis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
446.	Kalanchoe thyrsiflora (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
447.	Kalanchoe tomentosa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
448.	Kalonji <i>Nigella sativa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.

449.	Kalonji <i>Nigella sativa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
450.	Kalonji <i>Nigella sativa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
451.	Chilli Dry <i>Capsicum annum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
452.	Chilli Dry <i>Capsicum annum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
453.	Chilli Dry <i>Capsicum annum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
454.	Kasuri methi <i>Trigonella foenum-graecum</i>	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
455.	Kasuri methi <i>Trigonella foenum-graecum</i>	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
456.	Kasuri methi <i>Trigonella foenum-graecum</i>	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
457.	Ladies finger <i>Abelmoschus esculentus</i> , Pomegranate <i>Punica granatum</i> , Watermelon <i>Citrullus lanatus</i> , Garlic <i>Allium sativum</i> , Onion red <i>Allium cepa</i> , Onion white <i>Allium cepa</i> , Ginger <i>Zingiber Officinale</i> , Pineapple <i>Ananas comosus</i> , Lemon <i>Citrus limon</i> , Guava <i>Psidium guajava</i> , Turmeric <i>Curcuma longa</i> , Papaya <i>Carica papaya</i> , Beetroot <i>Beta vulgaris</i> , Beans Farm <i>Phaseolus vulgaris</i> , Banana Yelaki <i>Musa</i> , Brinjal Bottle <i>Solanum melongena</i> , Cucumber green <i>Cucumis sativus</i> , Cucumber French <i>Cucumis sativus</i> , Guava white <i>Psidium guajava</i> , Guava red <i>Psidium guajava</i> , Ginger <i>Zingiber officinale</i> , Lemon <i>Citrus limon</i> , Ladies finger <i>Abelmoschus esculentus</i> , Muskmelon <i>Cucumis melo</i> , Orange <i>Citrus sinensis</i> , Pineapple <i>Ananas comosus</i> , Papaya <i>Carica papaya</i> , Pomegranate <i>Punica granatum</i> , Tomato farm <i>Solanum lycopersicum</i> , Tomato natty <i>Solanum lycopersicum</i> , Turmeric <i>Curcuma longa</i> , Watermelon <i>Citrullus lanatus</i> , Green Grapes <i>Vitis vinifera</i> , Bangalore Blue Grapes <i>Vitis vinifera</i> , Baby Corn <i>Zea mays</i> , Corn <i>Zea mays</i> , , oconut <i>Cocos nucifera</i> , Beetroot <i>Beta</i>	Middle East (UAE, Qatar, Kuwait, Oman, Saudi Arabia)	Fresh Vegetables/Fruits/Flowers – UAE: The Consignment Is Free From Fall Army Worms (<i>Spodoptera frugiperda</i>) on The Basis Of Visual Inspection. (Chilli, Cucumber and Mango needs MRL Test Certificate.) Saudi Arabia: MRL Test Certificate No. for Chilli with inspection at accredited pack house. Other Countries no AD

	<p><i>vulgaris</i>, Yelaki banana <i>Musa</i>, Robusta banana <i>Musa</i>, Brinjal bottle <i>Solanum melongena</i>, French cucumber <i>Cucumis sativus</i>, Yam <i>Dioscorea alata</i>, Beans broad <i>Vicia faba</i>, Pumpkin red <i>Cucurbitan</i> Pumpkin white <i>Cucurbita</i>, Carrot ooty <i>Daucus carota subsp</i>, Sweet potato <i>Ipomoea batatas</i>, Potato <i>Solanum tuberosum</i>, Peas <i>Pisum sativum</i>, Jamun <i>Syzygium cumini</i>, Bitter Gourd <i>Momordica charantia</i>, Bottle gourd <i>Lagenaria siceraria</i>, Cucumber <i>Cucumis sativus</i>, Tindly <i>Coccinia grandis</i>, Long beans <i>Vicia</i>, Banana <i>Musa</i>, Drumstick <i>Moringa</i>, Sapota <i>Manilkara zapota</i>, Amla <i>Phyllanthus embilica</i>, Cluster beans <i>Cyamopsis tetragonoloba</i>, Yam <i>Dioscorea alata</i>, Colocassia <i>Colocasia esculenta</i>, Ridge Gourd <i>Luffa acutangula</i>, Mango <i>Mangifera indica</i>, Chilli <i>Capsicum annum</i>, Curry leaves <i>Murraya koenigii</i>, Chrysanthemum flowers <i>Chrysanthemum morifolium</i>, Jasmine <i>Jasminum</i>, Gerbera <i>Gerbera jamesonii</i>, Buttons <i>Centaurea cyanus</i>, Tuberose <i>Polianthes tuberosa</i>, Nerium <i>Nerium Oleander</i>, Gladiolus <i>Gladiolus sp.</i>, Rose <i>Rosa</i>, Kakada <i>Jasminum multiflorum</i>, Mary gold <i>Tagetes</i></p>		
458.	Lagenandra keralensis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
459.	Lagenandra keralensis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
460.	Lagenandra meeboldi 'Green'	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
461.	Lagenandra meeboldi 'Green (Tissue Culture)'	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
462.	Lagenandra meeboldi 'Red'	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
463.	Lagenandra meeboldi 'Red' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
464.	Lagenandra meeboldi 'Vinay chandra' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

465.	Lagenandra meeboldi 'Vinay chandra' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
466.	Lagenandra meeboldi 'silver queen' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
467.	Lagenandra meeboldii 'Green' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
468.	Lagenandra meeboldii 'Red' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
469.	Lagenandra meeboldii 'Vinay Chandra' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
470.	Lagenandra nairii (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
471.	Lenophyllum reflexum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
472.	Liliopesis novae zelandia (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
473.	Limnophila hippuroides (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
474.	Limnophila sessiliflora (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
475.	Lindernia anagallis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
476.	Liriope (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
477.	Liriope muscari Big Blue (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

478.	Liriope muscari Royal Purple (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
479.	Liriope muscari Samantha (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
480.	Liriope muscari (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
481.	Liriope muscari Evergreen Giant (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
482.	Liriope muscari(Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of Xylella fastidiosa as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IHR, Bangalore-Govt.of India
483.	Lomandra (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
484.	Lomandra confertifolia Wingarra(Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of Xylella fastidiosa as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IHR, Bangalore-Govt.of India

485.	Lomandra glauca Blue Ridge (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
486.	Lomandra longifolia (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
487.	Lomandra longifolia Lime Tuff (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
488.	Lomandra longifolia Little Con (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
489.	Lomandra longifolia Little Pal (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
490.	Lomandra longifolia(Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of Xylella fastidiosa as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IHR, Bangalore-Govt.of India
491.	Ludwigia arcuata (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
492.	Ludwigia arcuata 'Narrow leaf' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
493.	Ludwigia glandulosa (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

494.	Ludwigia inclinata (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
495.	Ludwigia ovalis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
496.	Ludwigia palustris 'Green' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
497.	Ludwigia repens 'super red' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
498.	Maize/ corn (<i>Zea mays</i>) Seed	Libya	Seeds were treated with thiram 2g per kg
499.	Maize/Corn (<i>Zea mays</i>)- Seed	Brazil	The seeds are treated with Captan @ gms per kg of seed
500.	Maize/Corn (<i>Zea mays</i>)- Seed	Indonesia	Seeds are free <i>Acremonium stricum</i> , <i>Fusarium sporotrichoides</i> , <i>Gloeocercospora sorghi</i> , <i>gaemannomyces graminis var graminis</i> , <i>Pernosclerospora sorghi</i> , <i>sclerophthora macrospora</i> , <i>Sphacelotheca reliana</i> , <i>Stenocarpella maydis</i> , <i>Sclerospora graminicola</i> , <i>Pernosclerospora philippinensis</i> , <i>Macrophomina phaseolina</i> , <i>Pantoea stewartii</i> , <i>Dickeya zea</i> , <i>Pantoea ananatis</i> , <i>Pseudomonas syringae pv. Syringae</i> , <i>Xanthomonas vasicola pv.Holcicola</i> , <i>Sitophilus granarium</i> , <i>Trogoderma granarium</i> , <i>Acnthoscelides obtus</i> , and certification that the seeds have undergone the routine procedure and found to be free from other seed borne plant Pathogensw storage insect pests and weed seeds
501.	Maize/Corn (<i>Zea mays</i>)- Seed	Mexico	Free from <i>Pseudomonas syringae pv. Lapsa Pseudomonas Psyringae pv. Syringae</i>
502.	Maize/Corn (<i>Zea mays</i>)- Seed	Pakistan	Certify that the seeds are tested and not infected by <i>Xanthomonas Stewart</i> Seeds were treated with chemical to protect the seeds from Fungi/Insects and the chemical used Thiram Applied at the rate of CC/Kg and Malathion Applied at the rate of CC/Kg
503.	Maize/Corn (<i>Zea mays</i>)- Seed	Philippines	The seeds are free from Aster Yellow <i>Phytoplasma</i> group <i>Pseudomonas syringae pv. Lapsa Pseudomonas psyringae pv. Syringae</i> <i>Acremonium maydis</i> , <i>Cochiliobolus carbonum</i> , <i>Fusarium sacchari</i> , <i>Sclerophthora rayssi ae var zea</i> , <i>Stenocarpella maydis</i> <i>Sclerophthora macrospora clerospora graminicola</i>
504.	Maize/Corn (<i>Zea mays</i>)- Seed	Tanzania	Import permit NO.Seeds were tested with captan 0.5 gm per kg and metalaxyl 4.0 gm per kg <i>Xanthomanas stewartii</i> Dowson is not known to occure in the country of production Maize lethel necrosis virus is not known to occure in the area of production
505.	Maize/Corn (<i>Zea mays</i>)- Seed	USA	NO AD
506.	Maize/Corn (<i>Zea mays</i>)- For consumption	Taiwan	The corn has been inspected and found free from the gumming disease (<i>Xanthomonas axonopodis pv. vasculorum</i>) (cobb) vauterin et.al
507.	Maize/Corn (<i>Zea mays</i>)- For consumption	Bangladesh	The plant and plant material are free from injurious insect pest and disease sand, soil. Merchandise shipped are throughly cleaned from soil and the plant and plant materials are free from injurious insect pest and disease or rendered so by effective traetment. Fumigation with methyl bromide (mbr) @ 32 grams/m3 for 24 hrs
508.	Maize/Corn (<i>Zea mays</i>)- For consumption	Reunion	The consignment is free from <i>Acrolepiopsis assectela</i> zeller, <i>Delia antique</i> and <i>Urocystis cepulae</i> , roots, leaves, soil, particle and or compost and other contaminants.

509.	Maize/Corn (<i>Zea mays</i>)- For consumption	Srilanka	Maize seeds are obtained from fields free from <i>Peronosclerospera hilippinenis</i> , <i>Cephalosporium maydis</i> weed striga, maize dwarf mosaic virus and <i>Eriwinia stewartii</i> or these pest do not occure in the country of production.
510.	Mango (<i>Mangifera indica</i>)- Fresh Fruit	UAE	The lot of mango accepted according to the codex alimantariuous maxium residue limits vide residue analysis report no.c2/0000011969 14.08.2018 issued by shriram institute for industrial research southern regional branch Bangalore karnataka India as per requirement of ministry of agriculture of UAE as certificate attached in original
511.	Mango (<i>Mangifera indica</i>)- Fresh Fruit	USA	<ol style="list-style-type: none"> 1. The mangoes were subjected to one of the pre- or post harvest mitigation options described in 7CFR 319.56-46.” and 2. The mangoes were inspected during preclearance activities and found free from <i>Cytosphaeramangiferae</i>, <i>Macrophomamangifera</i>, and <i>Xanthomonascampestrispv.mangiferaeindicae</i>.” 3. Import permit Number 4. The irradiation treatment has to be endorsed on the PSC. 5. OTHER REQUIREMWNTS: 6. Treatment by USDA-approved Irradiation treatment facility as per Standard Operating Procedure and Operational Work Plan. 7. Approved treatment for various commodities including Mango & Pomegranate from India is IR 400 Gy (T105-2-a)
512.	Mango (<i>Mangifera indica</i>)- Fresh Fruits	Australia	<p>The fruit in this consignment has been produced in India in accordance with the conditions governing entry of fresh mango fruit to Australia and in accordance with the Irradiation Operational Work Plan between India and Australia” Treatment Identification No.0752018a3001 Treated at Innova Bio Park Ltd, Malur (TFC-003/IR)</p> <p style="text-align: center;">Or</p> <ol style="list-style-type: none"> 1. a) “The mangoes in this consignment have been produced in India in accordance with the conditions governing entry of fresh mango fruit to Australia and inspected and found free of quarantine pests” AND b) “The mangoes in this consignment have been sourced from a designated place of production or production site in India which is free of <i>Sternochetus mangiferae</i> and <i>S. frigidus</i>” AND c) “The mangoes in this consignment have been sourced from a designated place of production or production site in India which is free of <i>Deanolis sublimbalis</i>”. Also the following information must be included on the Phytosanitary certificate:a) Treatment details, including date of treatment, temperature and duration of treatmentb) Packing house / treatment facility registration numbersc) Number of cartons per consignmentd) Container and seal number (seafreight only) 2. Vapour heat Treatment- a) Pulp core temperature of 46.5°C maintained for a minimum of 30 minutes; OR b) Pulp core temperature of 47.5°C maintained for a minimum of 20 minutes.

513.	Mango (<i>Mangifera indica</i>)- Fresh Fruits-	European Union Countries (Austria, Belgium, Bulgaria, Croatia, Cyprus, Czech, Denmark, Estonia, Finland, France, Germany, Greece, Hungary, Ireland, Italy, Latvia, Lithuania, Luxembourg, Malta, Netherland, Poland, Portugal, Romania, Slovakia, Slovenia, Spain, Sweden, UK)	<ol style="list-style-type: none"> 1. "The consignment of Mango treatment was supervised by(Name of PQ Station).... & found free from Tephritidae (Non-European) "Fulfills Annex I Part –A Section -I of EC Plant Health Directive 2000/29/EC as amended." 2. Vapor Heat Treatment- Pulp temperatures of the center of the fruits shall be 47.5°C and be kept continuously at 47.5°C or above for 30 minutes. Or 48°C and be kept continuously at 48°C or above for 20 minutes. or 3. Hot water immersion treatment- I- for 500 grams or below fruits – 48 °C water temperature for 60 min. II- for 501 grams to 750 grams fruits – 48 °C water temperature for 75 min. III- for 901 grams or above fruits – 48 °C water temperature for 90 min.
514.	Mango (<i>Mangifera indica</i>)- Fresh Fruits-	Switzerland	<ol style="list-style-type: none"> 1. "The consignment of Mango treatment was supervised by(Name of PQ Station).... & found free from Tephritidae (Non-European) "Fulfills Annex I Part –A Section -I of EC Plant Health Directive 2000/29/EC as amended." 2. Vapor Heat Treatment- Pulp temperatures of the center of the fruits shall be 47.5°C and be kept continuously at 47.5°C or above for 30 minutes. Or 48°C and be kept continuously at 48°C or above for 20 minutes. or 3. Hot water immersion treatment- I- for 500 grams or below fruits – 48 °C water temperature for 60 min. II- for 501 grams to 750 grams fruits – 48 °C water temperature for 75 min. III- for 901 grams or above fruits – 48 °C water temperature for 90 min.
515.	Mango (<i>Mangifera indica</i>)- Fresh Fruits-	New Zealand	<ol style="list-style-type: none"> 1. "The mangoes in this commercial treatment, consignment have (i) been visually inspected in accordance with appropriate official procedures and found free from regulated organisms and (ii) been produced in accordance with terms of, and treated as per Section 4.1 of the official assurance programme between MAF and Directorate of Plant Protection, Quarantine & Storage (PPQS), Ministry of Agriculture & Farmers Welfare, Government Of India." 2. Vapor Heat Treatment- Pulp temperatures of the center of the fruits shall be 48°C and be kept continuously at 48°C or above for 20 minutes. 3. Vapor heat treatment for export with saturated vapour in an approved vapour heat treatment facility in accordance with the program of Indian Plant Quarantine Authority. Fruit pulp temperature is raised until the fruit core reaches at least 48 0C for all sensors. The temperature is then maintained for at least 20 minutes in accordance with the National Standard for Phytosanitary Measures(NSPM 20).
516.	Mango (<i>Mangifera indica</i>)- Fresh Fruits-	Iran	The Mango fruit in this consignment has been produced in India in accordance with the conditions governing entry of fresh mango fruit to Iran and in accordance with the Hot water treatment plan between India and Iran. Hot water immersion treatment – 46.1-46.5 °C for 70 minutes.

			<p>Consignment was inspected and found free from any pathogenic symptoms and live infestation of following pests-</p> <table border="0"> <tr> <td>1. <i>Aleurodicus disperses</i></td> <td>2. <i>Aulacaspistubercularis</i></td> </tr> <tr> <td>3. <i>Bactrocera spp.</i></td> <td>4. <i>Ceroplastesrubens</i></td> </tr> <tr> <td>5. <i>Cryptoblabesgnidiella</i></td> <td>6. <i>Cytosphaeramangiferae</i></td> </tr> <tr> <td>7. <i>Dysmicoccusbrevipes</i></td> <td>8. <i>Elsinoemangiferae</i></td> </tr> <tr> <td>9. <i>Hendersoniacreberrima</i></td> <td>10. <i>Macrophomamangiferae</i></td> </tr> <tr> <td>11. <i>Nipaecoccusviridis</i></td> <td>12. <i>Phomopsismangiferae</i></td> </tr> <tr> <td>13. <i>Parasaissetianigra</i></td> <td>14. <i>Planococcuslilacinus</i></td> </tr> <tr> <td>15. <i>Pseudaulacaspiscockerlli</i></td> <td>16. <i>Pseudaonidiatrilobitiformis</i></td> </tr> <tr> <td>17. <i>Rastrococcusinvadens</i></td> <td>18. <i>Sternochetusfrigidus</i></td> </tr> <tr> <td>19. <i>Sternochetusmangiferae</i></td> <td>20. <i>Thripsalmi</i></td> </tr> <tr> <td>21. <i>Xanthomonascampestrispv.mangiferaeindicae</i></td> <td></td> </tr> </table>	1. <i>Aleurodicus disperses</i>	2. <i>Aulacaspistubercularis</i>	3. <i>Bactrocera spp.</i>	4. <i>Ceroplastesrubens</i>	5. <i>Cryptoblabesgnidiella</i>	6. <i>Cytosphaeramangiferae</i>	7. <i>Dysmicoccusbrevipes</i>	8. <i>Elsinoemangiferae</i>	9. <i>Hendersoniacreberrima</i>	10. <i>Macrophomamangiferae</i>	11. <i>Nipaecoccusviridis</i>	12. <i>Phomopsismangiferae</i>	13. <i>Parasaissetianigra</i>	14. <i>Planococcuslilacinus</i>	15. <i>Pseudaulacaspiscockerlli</i>	16. <i>Pseudaonidiatrilobitiformis</i>	17. <i>Rastrococcusinvadens</i>	18. <i>Sternochetusfrigidus</i>	19. <i>Sternochetusmangiferae</i>	20. <i>Thripsalmi</i>	21. <i>Xanthomonascampestrispv.mangiferaeindicae</i>	
1. <i>Aleurodicus disperses</i>	2. <i>Aulacaspistubercularis</i>																								
3. <i>Bactrocera spp.</i>	4. <i>Ceroplastesrubens</i>																								
5. <i>Cryptoblabesgnidiella</i>	6. <i>Cytosphaeramangiferae</i>																								
7. <i>Dysmicoccusbrevipes</i>	8. <i>Elsinoemangiferae</i>																								
9. <i>Hendersoniacreberrima</i>	10. <i>Macrophomamangiferae</i>																								
11. <i>Nipaecoccusviridis</i>	12. <i>Phomopsismangiferae</i>																								
13. <i>Parasaissetianigra</i>	14. <i>Planococcuslilacinus</i>																								
15. <i>Pseudaulacaspiscockerlli</i>	16. <i>Pseudaonidiatrilobitiformis</i>																								
17. <i>Rastrococcusinvadens</i>	18. <i>Sternochetusfrigidus</i>																								
19. <i>Sternochetusmangiferae</i>	20. <i>Thripsalmi</i>																								
21. <i>Xanthomonascampestrispv.mangiferaeindicae</i>																									
517.	Mango (<i>Mangifera indica</i>)- Fresh Fruits-	China	<p>(a) the mangoes inspected and found free from quarantine pests as specified in Annexure to Phytosanitary Protocol viz., (1) <i>Aonidiellacomperiei</i>; (2) <i>Bactroceracorrecta</i>; (3) <i>Bactroceraazonata</i>; (4) <i>Deanolisalbizonalis</i>; (5) <i>Eriosomyiaindica</i>; (6) <i>Parlatoriacrypta</i>, (7) <i>Sternochetusfrigidus</i>; (8) <i>Sternochetusmangiferae</i>; and (9) <i>Capnodiumramosum</i>; and</p> <p>(b) the mangoes covered by this phytosanitary certificate comply with the requirements established in the Phytosanitary Protocol on mangoes entry into China, which was duly signed between India and China on June 23, 2003.</p>																						
518.	Mango (<i>Mangifera indica</i>)- Fresh Fruits-	Japan	<ol style="list-style-type: none"> 1. THIS IS TO CERTIFY THAT THE MANGO FRUITS COVERED BY THIS CERTIFICATE ARE FREE FROM <i>BACTROCERA DORSALIS</i> SPECIES COMPLEX AND MELON FLY <i>BACTROCERA CUCURBITAE</i>. 2. Vapor Heat Treatment - In Vapor Heat Treatment facilities, the fruits shall be disinfested in saturated water vapor. The air temperature inside the treatment chamber shall be raised step by step to 50°C or above, so that the pulp temperatures of the center of the fruits shall be raised to 47.5°C and be kept continuously at 47.5°C or above for 20 minutes. 3. Pulp temperatures of the center of the fruits shall be 47.5°C and be kept continuously at 47.5°C or above for 20 minutes. 4. OTHER REQUIREMENTS: The consignment has to be treated in front of Japanese PQ officials. 																						
519.	Maranta leuconeura Kerchoveana (Tissue Culture)	Australia	<p>“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....</p>																						

520.	Maranta leuconeura Kerchoveana (Tissue Culture)	Netherlands/Holland	Certified that the consignment complies with point 32.1(d), 46 (b) (d) of Annex IV AI of Plant Health Directive (2000/29/EC)
521.	Maranta leuconeura Tricolor (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
522.	Maranta leuconeura Tricolor (Tissue Culture)	Netherlands/Holland	Certified that the consignment complies with point 32.1(d), 46 (b) (d) of Annex IV AI of Plant Health Directive (2000/29/EC)
523.	Maranta leuconeura (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
524.	Maranta leuconeura Blue Marxil (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
525.	Mayaca fluviatilis (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
526.	Melaleuca thymifolia White Lace (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
527.	Metrosideros tementosa (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
528.	Metrosideros tementosa Fiji Fire (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
529.	Metrosideros tementosa Spring Fire (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this

			consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
530.	Micranthemum sp (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
531.	Micranthemum sp. (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
532.	Micranthemum umbrosum (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
533.	Microsorium pteropus 'Narrow leaf' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
534.	Microsorium pteropus 'malabar' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
535.	Microsorium pteropus 'Trident'	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
536.	Microsorium pteropus 'Trident (Tissue Culture)'	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
537.	Microsorium sp. 'Trident' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
538.	Milingtonia hortensis (Tissue Culture)	UAE	"The consignment is free from Fall Army Worm (<i>Spodoptera frugiperda</i>) on the basis of visual inspection".
539.	Monstera adonsonii (Tissue Culture)	Australia	"Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened". AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
540.	Monstera deliciosa (Tissue Culture)	Australia	"Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened". AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
541.	Monstera deliciosa Variegated (Tissue Culture)	Australia	"Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened". AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

542.	Monstera obliqua (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
543.	Monstera tauerii (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
544.	Monstera tauerii Variegated (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
545.	Mung <i>Vigna radiate</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
546.	Mung <i>Vigna radiate</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
547.	Mung <i>Vigna radiate</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
548.	Murraya paniculata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
549.	Musa basjoo (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
550.	Musa basjoo(Tissue Culture)	Germany	Certified that the live tissue culture plants (Musa basjoo) in this consignment were tested and found to be free from bacterium <i>Ralstonia solanacearum</i> and fulfils item 25.7 (b) of Annex IV AI of Plant Health Directive (2000/29/EC) as amended
551.	Musa ensete (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

552.	Musa grand nain (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
553.	Musa spp(Tissue Culture)	Angola	Certified that plantlets were raised through sterile vtissue culture procedure and free of virus including Banana Bunchy Top Virus, Cucumber Mosaic Virus, Banana Streak Virus, Banana Mosaic Virus, Banana Bract Mosaic Virus, Xanthomonas (Banana Wilt) using ELIZA/PCR test, Erwinia Chrysathemi, Pseudomos Syringae Pv Syringae (Bacterial Canker)
554.	Musa spp(Tissue Culture)	Ethiopia	The plants are certified plants and vius indexed and free of virus and other diseases and pests, and free from natural soil
555.	Musk Melon (Cucumis melo)- Seed	Jordan	Nil
556.	Musk Melon (Cucumis melo)- Seed	South Korea	Nil
557.	Myriophyllum hippuroides (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
558.	Mysore Mamra (South Indian Mamra) <i>Oryza sativa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
559.	Mysore Mamra (South Indian Mamra) <i>Oryza sativa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
560.	Mysore Mamra (South Indian Mamra) <i>Oryza sativa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
561.	Nasaea pedicilata (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
562.	Neomarica caerulea (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
563.	Neomarica caerulea Indigo Blue (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

564.	Neomarica fluminensis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
565.	Neomarica gracilis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
566.	Neomarica longifolia Yellow (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
567.	Neomarica northiana (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
568.	Neomarica regida (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
569.	Neomarica sabinei Alba (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
570.	Okra (<i>Abelmoschus esculentus</i>) Seed	Bangladesh	Seeds are free from Quarantine weed seeds
571.	Okra (<i>Abelmoschus esculentus</i>) Seed	Libya	Seeds were treated with thiram 2g per kg
572.	Okra (<i>Abelmoschus esculentus</i>) Seed	Malaysia	Seeds are free following Pests Weed: i. <i>Parthenium hysterophorus</i>
573.	Okra (<i>Abelmoschus esculentus</i>) Seed	Nepal	Consignment of Plants and Plant Products is free from soil, seeds of weeds and other infection.

574.	Okra (<i>Abelmoschus esculentus</i>) Seed	Sudan	Seeds are free from soil and weed seeds including quarantine weed seeds Free from <i>ralstonia solanacerum</i> , <i>clavibacter michiganens</i> , <i>pseudomonas</i> sps and <i>phytophthera</i> sps. Treated with fungicide and streptomycin and red polymer
575.	Okra (<i>Abelmoschus esculentus</i>) Seed	Tanzania	The seeds were harvested from mother plants which were inspected during active growth and found free from pests.
576.	Okra (<i>Abelmoschus esculentus</i>) Seed	USA	Nil
577.	Onion (<i>Allium cepa</i>)- Seed	Tanzania	Nil
578.	Onion (<i>Allium cepa</i>)- Bulb for consumption	Mauritius	This consignment is free from <i>Urocystis cepulae</i> , <i>Delia antiqua</i> , <i>Aceria tulipae</i> , thrips, stem and bulb nematode (<i>Ditylenchus dipsaci</i>) fumigation with methyl bromide (mbr) @ 16 grams/m ³ for 12 hrs
579.	Onion (<i>Allium cepa</i>)- Bulb for consumption	Taiwan	Consignment is free from mites, <i>Rhizoglyphus echinopus</i> , <i>Ditylenchus dipsaci</i> & free from roots, soil, particle or compost.
580.	Onion (<i>Allium cepa</i>)- Bulb for consumption	Indonesia	The consignment is free from roots, leaves, soil particle and or compost. The products are free from contamination of plant debris, soil and other inert matter.
581.	Ophiopogon planiscapus Black Mondo (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
582.	Ophiopogon planiscapus Large Mondo (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
583.	Ophiopogon planiscapus Stripey White (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
584.	Oxalis triangularis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

585.	<i>Oxalis tubistipula</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
586.	<i>Pachyphytum oviferum</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
587.	<i>Pellonia repens</i> (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
588.	<i>Peperomia argyreia</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
589.	<i>Peperomia asperula</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
590.	<i>Peperomia caperata</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
591.	<i>Peperomia clusiifolia</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
592.	<i>Peperomia dolabriformis</i> (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

593.	Peperomia magnoliifolia (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
594.	Peperomia magnoliifolia(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
595.	Peperomia obtusifolia Green (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
596.	Peperomia obtusifolia Variegated (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
597.	Petunia (Petunia sp.)-Seed	Czech republic	Nil
598.	Philodendron (<i>Philodendron</i> sp.)- Tissue Culture plants	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found free from any symptoms of diseases or microbial infection. Prior to removal of the plant tissue from the media, the tissue cultures were inspected and found to be free of contamination. The plant tissues were aseptically transferred, under supervision, to sterile containers and sealed, and not subsequently re- opened.All tissue cultures in this consignment were derived from mother tissue cultures that were tested by polymerase chain reaction (PCR) and found free of <i>Xylella fastidiosa</i> as indicated on laboratory test number __dated: These plants are produced at their AQIS accredited lab at _____
599.	Philodendron (<i>Philodendron</i> sp.)- Tissue Culture plants	USA	Tissue Cultures in this consignment has been grown In Vitro and produced within a secure environment in a production area that is free of Potato Cyst Nematodes and has never been grown in soil nor come in contact with soil
600.	Philodendron (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

601.	Philodendron cordatum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
602.	Philodendron eximium (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
603.	Philodendron hastatum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
604.	Philodendron mamei Silver Cloud (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
605.	Philodendron scanden Brasil (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
606.	Philodendron selloum Maccollys Finale (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
607.	Philodendron selloum (Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of Xylella fastidiosa as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IHR, Bangalore-Govt.of India

608.	Philodendron selloum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
609.	Philodendron selloum Birkin (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
610.	Philodendron selloum Black Cardinal (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
611.	Philodendron selloum Burle Marx (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
612.	Philodendron selloum Celon Beauty (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
613.	Philodendron selloum Congo (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
614.	Philodendron selloum Goeldii (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

615.	Philodendron selloum Goeldii(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
616.	Philodendron selloum Hope (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
617.	Philodendron selloum Imperial Gold (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
618.	Philodendron selloum Moonshine (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
619.	Philodendron selloum Pink Magic (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
620.	Philodendron selloum Pink Princes (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
621.	Philodendron selloum Prince of Orange (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

622.	Philodendron selloum Rojo Congo (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
623.	Philodendron selloum Silver Sword (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
624.	Philodendron selloum Super Atom (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
625.	Philodendron selloum White Blizzard (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
626.	Philodendron selloum Xanadu (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
627.	Philodendron selloum(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
628.	Philodendron selloum(Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
629.	Philodendron sp. 'Papua New Guinea' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

630.	Philodendron spp (Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of Xylella fastidiosa as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IHR, Bangalore-Govt.of India
631.	Philodendron spp(Tissue Culture)	USA	Tissue cultures in this consignment has been grown <i>InVitro</i> , and produced within a secure environment in a production area that is free of Potato cyst nematodes and has never been grown in soil nor come in contact with soil, as indicated on laboratory test dated XXXXX29/01/2019 IHR, Bangalore-Govt.of India.
632.	Philodendron squamiferum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
633.	Philodendron xanadu(Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of Xylella fastidiosa as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IHR, Bangalore-Govt.of India
634.	Phormium (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
635.	Pilea peperomioides (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
636.	Pitcairnia sanguinea (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

637.	Pitcairnia sanguinea(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (<i>Globodera rostochiensis</i>) and (<i>Globodera pallida</i>)”
638.	Pogostemon erectus (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
639.	Polygonum sp. 'Pink' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
640.	Potato, <i>Solanum tuberosum</i>	Mauritius	This consignment is free from <i>Ralstonia solanacearum</i> , <i>Clavbacter michiganesis</i> sub. <i>Sepedonicum</i> , <i>Synchytrium endobioticum</i> , potato cyst nematode, <i>Spongospora subterranea</i> , colorado beetel, <i>Leptinotarsa decemline</i>
641.	Proiphys cunninghamii (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
642.	Proiphys infundibularis (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
643.	Pumpkin (<i>Cucurbita moschata</i>)- Seeds	Japan	"The parent plants are grown from seeds disinfected against <i>Acidovorax avenae</i> subsp. <i>citrulli</i> or known to be free from this pest AND The parent plants and fruits at a place of production or a production site (including a plant growth facility) are found to be free from <i>Acidovorax avenae</i> subsp. <i>citrulli</i> by inspection, including laboratory testing of any suspicious symptoms, carried out during fruit maturity stage before harvesting".
644.	Pumpkin (<i>Cucurbita moschata</i>)- Seeds	Korea	NIL
645.	Pumpkin (<i>Cucurbita moschata</i>)- Seeds	Philippines	Seeds are free from soil
646.	Pumpkin (<i>Cucurbita moschata</i>)- Seeds	Philippines	Seeds free from the soil
647.	Ragi , <i>Eleusine coracana</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
648.	Ragi , <i>Eleusine coracana</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
649.	Ragi , <i>Eleusine coracana</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
650.	Ranunculus inundates (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

651.	Chilli Dry <i>Capsicum annum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
652.	Chilli Dry, <i>Capsicum annum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
653.	Chilli Dry, <i>Capsicum annum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
654.	Riccardia chamedryfolia (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
655.	Rice Flour Buen <i>Oryza sativa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
656.	Rice Flour Buen <i>Oryza sativa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
657.	Rice Flour Buen <i>Oryza sativa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
658.	Rice Flour Coarse <i>Oryza sativa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
659.	Rice Flour Coarse <i>Oryza sativa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
660.	Rice Flour Coarse <i>Oryza sativa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
661.	Rice Flour Udupi <i>Oryza sativa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
662.	Rice Flour Udupi <i>Oryza sativa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
663.	Rice Flour Udupi <i>Oryza sativa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
664.	Rice Flour Udupi <i>Oryza sativa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
665.	Rice Flour Udupi <i>Oryza sativa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
666.	Rice Flour Udupi <i>Oryza sativa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
667.	Rice <i>Oryza sativa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
668.	Rice <i>Oryza sativa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.

669.	Rice <i>Oryza sativa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
670.	Roasted Gram <i>Cicer arietinum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
671.	Roasted Gram <i>Cicer arietinum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
672.	Rose (<i>Rosa</i> sp.)- Cut Flowers	Lebanon	"The produce is free of <i>Xylella Fastidiosa</i> and <i>Xylella Fastidiosa</i> is not known to occur in INDIA."
673.	Rose (<i>Rosa</i> sp.)- Cut Flowers	New Zealand	Rosa spp. have been treated by being immersed up to 50mm below the flowers in Glyphosate solution @ 15ml / 985ml of water of 20min. Temp. During the devitalization Process was 28-32°C. It was inspected in accordance with appropriate procedure & found to be free of regulated organisms specified by NZ MAF
674.	Rose (<i>Rosa</i> spp.) Cut flowers	European Union	Fulfils item 45.2, option b of Annex IV AI of Plant Health Directive (2000/29/EC) as amended. Immediately prior to their export, they have been officially inspected and found free from <i>Bemisia tabaci</i> Genn. (Non-European populations).
675.	Rose (<i>Rosa</i> spp.)- Cut Flowers	Australia	Produced and prepared for export under approved system approachProduced has packed in pest proof numbered cartons that eliminate the possibility of entry or egress of insects pest.The strength of glyphosate used is 360g/l & mixed at the rate of 15ml/ 985ml water, rose stems were immersed for 20min to within 5cm of the head of for at least 35 cm depth. Also the temp. Of the roses at the time of dipping was 20deg.cent. All the flowers have been treated with glyphosate to render the buds non-viable and devitalisation done under our supervision at _____ltd.
676.	Rose (<i>Rosa</i> spp.)-Cut Flowers-	Malaysia	This consignment is free from fungi <i>Peronospora sparsa</i> (Downy mildew). Fungicide + Insecticide treatment. Import permit No.
677.	Rotala macrandra 'Bangladesh' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
678.	Rotala macrandra 'Pink' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
679.	Rotala macrandra 'Red' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
680.	Rotala nanjean (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
681.	Rotala rotundifolia (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
682.	Rotala rotundifolia colorata (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
683.	Rotala rotundifolia 'Fujian' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
684.	Rotala rotundifolia green (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
685.	Rotala sp 'Bangladesh' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

686.	Rotala sp 'Birunani' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
687.	Rotala sp 'Bonsai' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
688.	Rotala sp 'Manipur' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
689.	Rotala sp. 'Ceylon' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
690.	Rotala sp. 'Hra' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
691.	Rotala sp. 'Wayanad' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
692.	Rotala wallichii (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
693.	Sansevieria trifasciata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
694.	Scindapsus pictus (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
695.	Scrophulariaceae limnophila sp. (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
696.	Sesame seed <i>Sesamum indicum</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
697.	Sesame seed <i>Sesamum indicum</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
698.	Sesame seed <i>Sesamum indicum</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
699.	Jeera, <i>Elwendia persica</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
700.	Jeera <i>Elwendia persica</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
701.	Jeera <i>Elwendia persica</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.

702.	Sinningia bullatta (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
703.	Sonerila heterostemon (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
704.	Sonerila mamei (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
705.	Spathiphyllum floribundum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
706.	Spathiphyllum floribundum Blue Moon (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
707.	Spathiphyllum floribundum Loretta (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
708.	Spathiphyllum floribundum Sensation (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
709.	Spathoglottis plicata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
710.	Spathyphyllum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of

			Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
711.	Spathyphyllum (Tissue Culture)	USA	Tissue cultures in this consignment has been grown <i>InVitro</i> , and produced within a secure environment in a production area that is free of Potato cyst nematodes and has never been grown in soil nor come in contact with soil, as indicated on laboratory test dated XXXXX29/01/2019 IIHR, Bangalore-Govt.of India.
712.	Sponge guard (Luffa aegyptiaca)- seed	Taiwan	Seeds had been thoroughly inspected and found free from spindle tuber viroid based on official laboratory anaysis
713.	Sponge guard (Luffa aegyptiaca)- seed	Vietnam	NO AD
714.	Squash (<i>Cucurbita maxima</i>)- Seed	Japan	“Fulfills item 19 of the Annexed Table 2-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No73/1950)”. The parent plants are grown from seeds disinfected against this pest or known to be free from this pest And The parent plants and fruits at a place of production or a production site (including a plant growth facility) are found to be free from Acidovorax avenae subsp. citrulli by inspection, including laboratory testing of any suspicious symptoms, carried out during fruit maturity stage before harvesting. The parent plants were officially inspected during the active growth and found free from Khapra Beetle (Trogoderma Spp.), Bacterial Spot (Xanthomonas campestris pv. cucurbita), Bacterial Wilt (Erwinia tracheiphila), SCAB (Cladosporium cucumerinum), Angular Leaf Spot, Root Rot, Gummy Stem Blight, Musk Melon Mosaic Virus, Cucumber Green Mottle Mosaic Virus, Squash Mosaic Virus, Zucchini Yellow Mosaic Virus(ZYMV), Tobacco ringspot nepovirus, Tomato ringspot nepovirus, and Watermelon Bacterial Fruit Blotch. Free from Orobancha spp., Striga spp..
715.	Squash (<i>Cucurbita maxima</i>)- Seed	South Korea	Nil
716.	Syngonium (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
717.	Syngonium podophyllum (Tissue Culture)	Australia	Tissue cultures in this consignment were visually inspected immediately prior to export and found to be free from any symptoms of disease or microbial infection” “Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. All tissue cultures in this consignment were produced from mother tissue culture plants that were tested by PCR and found free of Xylella fastidiosa as indicated on laboratory test number XXXXX (TCPT001/19) dated XXX IIHR, Bangalore-Govt.of India

718.	Syngonium podophyllum (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
719.	Syngonium podophyllum Bronze (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
720.	Syngonium podophyllum Green Pixie (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
721.	Syngonium podophyllum Neon (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
722.	Syngonium podophyllum Pink Vein (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
723.	Syngonium podophyllum PPN (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
724.	Syngonium podophyllum PPN(Tissue Culture)	USA	“Certified that the live tissue culture plants in this consignment were produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (Globodera rostochiensis) and (Globodera pallida) ”
725.	Syngonium podophyllum Red Vein (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

726.	Syngonium podophyllum Silver Pearl (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
727.	Syngonium podophyllum White Butterfly (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
728.	Syngonium spp(Tissue Culture)	USA	Tissue cultures in this consignment has been grown InVitro, and produced within a secure environment in a production area that is free of Potato cyst nematodes and has never been grown in soil nor come in contact with soil, as indicated on laboratory test dated XXXXXXXXXX IIHR, Bangalore-Govt. of India
729.	Syzygium australe (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
730.	Tacca chantrieri (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
731.	Tacca integrifolia (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
732.	Tamarind Slabs <i>Tamarindus indica</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
733.	Tamarind Slabs <i>Tamarindus indica</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
734.	Tamarind Slabs <i>Tamarindus indica</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (<i>Trogoderma granarium</i>) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
735.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
736.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)

737.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
738.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
739.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
740.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
741.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
742.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
743.	Taxiphyllum barbieri (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
744.	Taxiphyllum sp. 'Flame moss' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
745.	Taxiphyllum sp. 'Flame moss' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
746.	Taxiphyllum sp. 'Peacock moss'	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
747.	Taxiphyllum sp. 'Peacock moss' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
748.	Taxiphyllum sp. 'Peacock moss' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
749.	Taxiphyllum sp. 'Spiky moss' (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
750.	Thespesia populnea (Tissue Culture)	UAE	"The consignment is free from Fall Army Worm (<i>Spodoptera frugiperda</i>) on the basis of visual inspection".
751.	Tissue culture plants	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
752.	Tissue culture plants	Kenya	Certified that the live tissue culture plants in this consignment were propagated and grown in sterile medium
753.	Tissue culture plants	USA	Certified that the live tissue culture plants in this consignment produced from mother tissue culture plants that were tested within 12 months preceding issuance of the Phytosanitary certificate and found free from Potato cyst nematodes (<i>Globodera rostochiensis</i>) and (<i>Globodera pallida</i>)
754.	Tomato (including <i>Lycopersicon esculentum</i> (= <i>Solanum lycopersicum</i>), <i>S. arcanum</i> , <i>S. cheesmaniae</i> , <i>S. chilense</i> , <i>S. galapagense</i> , <i>S. peruvianum</i> , <i>S. pimpinellifolium</i>)- Fresh fruit	Japan	Fulfils item 2 of the Annexed Table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No73/1950)

755.	Tomato (<i>Solanum lycopersicum</i>) Seed	Bangladesh	Plant and Plant materials are free from injurious insect pests and diseases Seeds are free from Bacterial Cankers (<i>Clavibacter michiganensis</i>), Bacterial Leaf spot (<i>Pseudomonas syringe</i> pv.tomato) Bacterial Pustule (<i>Pseudomonas syringe</i> pv.punctulens) Potato spindle tuber viroid, <i>Peronosporahyacyamipv.tabacina</i> , <i>Phoma andigena</i> , <i>Verticillium alboatrum</i> , <i>Clavibactermic higanensissubsp.sepodonicus</i> , Pepino Mosaic Virus, Tomato black ring Virus , Tomato bushy stunt virus, Tomato ring spot virus Free from quarantine weed seed.
756.	Tomato (<i>Solanum lycopersicum</i>) Seed	European Union	Consignment complies with Annex. IV. A. I Point 48 B of EC Plant Health Directive 2000/29/EC Consignment complies with Annex. I. A. of EC Plant Health Directive 2000/200/EC Hybrid tomato seeds were harvested from the parent plants inspected during active growth and found free from pseudomonas syreingae, Tomato ring spot virus, black ring virus, tomato spotted wilt virus and verticilium albo atrum.
757.	Tomato (<i>Solanum lycopersicum</i>) Seed	Jamaica	Seeds are free from Raistonia solanacearum and <i>Clavibacter michiganensis</i> spedonicus ,
758.	Tomato (<i>Solanum lycopersicum</i>) Seed	Jordan	Nil
759.	Tomato (<i>Solanum lycopersicum</i>) Seed	Kenya	The Seed was harvested from fields which have been inspected, during active growth and found to be free from <i>Didymella lycopersici</i> Kleb(Syn. <i>Ascochyta lycopersici</i> (Plowr) Brunaud, <i>Sphaeronaema lycopersici</i> Plowr, <i>Phoma lycopersici</i> Cooke) <i>Pseudomonas syringae</i> pv.tomato, Alfalfa mosaic virus, Cucumber mosaic virus, Tomato mosaic virus, Tobacco mosaic virus, Columnea latent viroid (CLVd) and Potato Spindle tuber viroid. Seeds treated with Polymer + Thiram + Thiamethoxam
760.	Tomato (<i>Solanum lycopersicum</i>) Seed	Malaysia	The seeds are free from the following Pests: Insect: Tuta absolute (tomato leaf minor), Virus:Tobacco Streak Virus, Tomato Ring Sport Virous, Arabis Mosaic , Weed: <i>Parthenium hysterophorus</i>
761.	Tomato (<i>Solanum lycopersicum</i>) Seed	Mauritius	(<i>Solanum lycopersicum</i>) Free from (a) <i>Pseudomonas syringae</i> pv tomato (b) <i>Xanthomonas Vesicatoria</i> (c) <i>Clavibacter michiganensis</i> subsp michiganensis (d) Potato spindle tuber viroid (PSTVd) (e) Tomato bushy stunt virus (TBSV) (f) Pepino mosaic virus (PepMV) (g) Tomato ringspot virus (ToRSV) (h) Tomato black ring virus(TBRV)
762.	Tomato (<i>Solanum lycopersicum</i>) Seed	Republic of Korea	Nil
763.	Tomato (<i>Solanum lycopersicum</i>) Seed	Sudan	Seeds are free from soil and weed seeds including quarantine weed seeds Free from ralstonia solanacerum, <i>clavibacter michiganens</i> , <i>pseudomonas</i> sps and <i>phytophthera</i> sps. Treated with fungicide and streptocycline and red polymer
764.	Tomato (<i>Solanum lycopersicum</i>) Seed	Taiwan	The seeds had been thoroughly inspected and found free from Potato spindle tuber viroid based on official laboratory analysis.
765.	Tomato (<i>Solanum lycopersicum</i>) Seed	Tanzania	The seeds harersted from plants which were inspected during active growth and found free from <i>Didymella lycopersici</i> , Tomato yellow leaf curl virus and Phtophthora sp.
766.	Tomato (<i>Solanum lycopersicum</i>) Seed	Tanzania	Nil
767.	Tomato (<i>Solanum lycopersicum</i>) Seed	Turkey	Seeds of <i>Lycopersicum esculentum</i> Mill. (tomato) the seeds have been obtained by meansof an appropriate acid extraction method and the field was inspected and found to be freefrom <i>Clavibacter michiganensis</i> subsp. michiganensis, <i>Xanthomonas vesicatoria</i> and PotatoSpindle tuber pospiviroid are not known to occur.

768.	Tomato (<i>Solanum lycopersicum</i>) Seed	Vietnam	Seeds were treated with fungicides
769.	Tomato (<i>Solanum lycopersicum</i>)- Seed	Bulgaria	Nil
770.	Tomato (<i>Solanum lycopersicum</i>)- Seed	Egypt	The consignment was tested and found free from the following seed borne virus: Tomato mosaic virus Tomato ring spot virus, Tomato yellow leaf curl virus
771.	Tomato (<i>Solanum lycopersicum</i>)- Seed	Jordan	Nil
772.	Tomato (<i>Solanum lycopersicum</i>)- Seed	Taiwan	Seeds had been thoroughly inspected and found free from spindle tuber viroid based on official laboratory analysis
773.	Tomato (<i>Solanum lycopersicum</i>)- Seed	USA	Seeds were found free from <i>Clavibacter michiganensis</i> ssp. <i>Michiganensis</i> , <i>Xanthomonas</i> (<i>campestris</i> pv.) <i>vesicatoria</i> , Potato spindle tuber viroid, Pepino mosaic virus Seeds treated with Polymer and Thiram
774.	Tomato (<i>Solanum lycopersicum</i>)- Seed	USA	Seeds were tested and found to be free from: Bacteria: <i>Acidovorax citrulli</i> , <i>Candidatus Phytoplasma solani</i> , <i>Corynebacterium michiganensis</i> <i>Corynebacterium sepedonicum</i> , <i>Dickeya dadantii</i> , <i>Clavibacter michiganensis</i> subsp. <i>michiganensis</i> , <i>Erwinia carotovora</i> spp. <i>Carotovora</i> , <i>Pseudomonas</i> spp., <i>Pseudomonas corrugate</i> , <i>Pseudomonas cichorii</i> , <i>Pseudomonas syringae</i> , <i>Pseudomonas syringae</i> pv. <i>tomato</i> , <i>Pseudomonas syringae</i> pv. <i>Punctulens</i> , <i>Pseudomonas syringae</i> pv. <i>Syringae</i> , <i>Ralstonia solanacearum</i> (syn. <i>Pseudomonas solanacearum</i>), <i>Pseudomonas viridiflava</i> , <i>Pseudomonas savastanoi</i> pv. <i>Phaseolicola</i> , <i>Xanthomonas campestris</i> pv. <i>vesicatoria</i> , <i>Xanthomonas campestris</i> pv. <i>campestris</i> , <i>Xanthomonas campestris</i> pv. <i>Perforans</i> , <i>Xanthomonas campestris</i> pv. <i>Gardneri</i> , <i>Xanthomonas campestris</i> pv. <i>Euvesicatoria</i> . Fungi: <i>Boeremia lycopersici</i> (syn. <i>Didymella lycopersici</i>), <i>Chalara elegans</i> , <i>Fusarium</i> spp <i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> , <i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> , <i>Gibberella</i> spp., <i>Peronospora hyoscyami</i> f. sp. <i>tabacina</i> , <i>Phoma andigena</i> , <i>Phoma destructive</i> , <i>Phytophthora</i> spp., <i>Phytophthora capsici</i> ., <i>Phytophthora cryptogea</i> ., <i>Phytophthora infestans</i> , <i>Puccinia pittieriana</i> (syn. <i>Gerwasia pittieriana</i>)., <i>Pythium</i> spp., <i>Sclerotinia sclerotiorum</i> , <i>Spongospora subterranea</i> f. sp. <i>Subterranean</i> , <i>Synchytrium endobioticum</i> , <i>Thecaphora solani</i> (<i>Angiosorus solani</i>), <i>Verticillium albo-atrum</i> , <i>Verticillium dahlia</i> <i>Verticillium tricorpus</i> . Viruses: Alfalfa mosaic virus (AMV), Arabis mosaic virus (ArMV) , Broad bean wilt virus (BBWV), Columnea latent viroid (CLVd), Cucumber mosaic virus (CMV), Peanut stunt virus (PSV) Pelargonium zonate spot ourmiavirus (PZSV), Pepino mosaic virus (PepMV), Pepper chat fruit viroid (PCFVd) Pepper mild mottle virus (PMMoV), Potato mop top virus (PMTV), Potato spindle tuber viroid (PSTVd) Tobamovirus spp., Tobacco mosaic virus (TMV), Tobacco rattle virus (TRV), Tobacco ringspot virus (TRSV), Tobacco streak virus (TSV), Tobacco mild green mosaic virus (TMGMV), Tomato apical stunt viroid (TASVd), Tomato aspermy virus (TAV), Tomato chlorotic dwarf viroid (TCDVd), Tomato black ring nepovirus (TBRV), Tomato bunchy top virus (TBTV), Tomato bushy stunt virus (TBSV), Tomato chlorotic dwarf viroid (TCDVd), Tomato infectious chlorosis virus (TICV), Tomato mottle mosaic virus (ToMMV), Tomato mosaic virus (ToMV), Tomato planta macho viroid (TPMVd), Tomato ringspot virus (ToRSV), Tomato spotted wilt virus (TSWV), Tomato yellow leaf curl virus (TYLCV), Tomato brown rugose fruit virus (ToBRFV). Nematodes: <i>Aphelenchoides ritzemabosi</i> , <i>Ditylenchus destructor</i> , <i>Ditylenchus dipsaci</i> , <i>Globodera pallida</i> . <i>Globodera rostochiensis</i> , <i>Longidorus elongates</i> , <i>Longidorus micoletzky</i> , <i>Meloidogyne</i> spp., <i>Meloidogyne goeldi</i> , <i>Paratrichodorus</i> spp., <i>Pratylenchus</i> spp., <i>Radopholus similis</i> , <i>Trichodorus</i> spp., <i>Xiphinema americanum</i> , <i>Xiphinema</i>

			diversicaudatum, Xiphinema index, Xiphinema rivesi Dalmasso. Weeds: Agropyron repens, Amaranthus retroflexus, Cassytha filiformis, Commelina benghatensis, Convolvulus arvensis, Crupina vulgaris, Cuscuta spp., Imperata cylindrical, Ipomoea aquatic, Lolium temulentum L., Orobanche spp., Oryza rufipogon, Solanum rostratum, Sonchus arvensis, Striga spp. Insects: Aphis craccivora, Dermestidae spp., Frankliniella occidentalis, Leptinotarsa decemlineata, Liriomyza trifolii, Mythimna unipuncta, Spodoptera littoralis, Tenebrio molitor, Thrips tabaci, Trogoderma spp., Trogoderma granarium, Trogoderma variabile
775.	Tulbaghia spp (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
776.	Tulbaghia spp(Tissue Culture)	United Kingdom	Certified that the consignment complies with Annex IV.A.I., point 32.1(d), 32.3 (d), 36.1(d), 45.1 (d) and point 46 (b) option (d) of EC Plant Health Directive 2000/29/EC
777.	Tulbaghia violacea Purple Eye (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
778.	Tulsi (<i>Ocimum tenuiflorum</i>), Leafy Vegetables-	European Union	Fulfils item 45.2 option b of the annex IV AI of plant health directive (2000/29/EC) as amended. Immediately prior to their export, the leafy vegetables have been officially inspected and found free from Bemisia tabaci genn (Non-European population). Fulfils item 32.2 option b of the annex IV AI of plant health directive (2000/29/EC) as amended. Immediately prior to their export, the leafy vegetables have been officially inspected and found free from <i>Liriomyza sativae</i> (Blanchard) and <i>Amauromyza maculosa</i> (Malloch).
779.	Turmeric <i>Curumma longa</i> (Food Product)	Australia	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
780.	Turmeric <i>Curumma longa</i> (Food Product)	Canada	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
781.	Turmeric <i>Curumma longa</i> (Food Product)	USA	This shipment has been inspected and found free of khapra beetle (Trogoderma granarium) Treatment: Fumigation with MBr at 80g/m3 for 48 hours at 21 degree celcius.
782.	Vesicularia montagnei (Tissue Culture)	Japan	Fulfils item 7 of the Annexed table 1-2 of the Ordinance for Enforcement of the Plant Protection Act (MAF Ordinance No.73/1950)
783.	Watermelon (<i>Citrullus lanatus</i>)- Seed	Bulgaria	Nil
784.	Watermelon (<i>Citrullus lanatus</i>)- Seed	Jamaica	Seeds are free from Didymella bryoniae, Pseudomonas syringae pv.lachrymans and Xathomonas campstris pv cucurbitae.

785.	Watermelon (<i>Citrullus lanatus</i>)- Seed	Kenya	The seeds was harvested from plants that have been inspected during active growth and found to be free from cucumber mosaic virus, tomato spot virus, acidovorax citrull, alfalfa mosaic virus, squash mosaic virus, zucchini yellow mosaic virus, Seeds treated with Polymer + Thiram
786.	Watermelon (<i>Citrullus lanatus</i>)- Seed	New Zealand	The <i>Citrullus lanatus</i> seeds for sowing in this consignment have been: sourced from a seed lot officially sampled according to ISTA or AOSA methodology, and tested using the ISTA validated ELISA or a NPPO approved PCR method and found free from Cucumber green mottle mosaic virus. The <i>Citrullus lanatus</i> seeds for sowing in this consignment have been: sourced from a seed lot officially sampled according to ISTA or AOSA methodology, and tested using a NPPO approved serological (ELISA) or molecular (PCR) method and found free from Kyuri green mottle mosaic virus.
787.	Watermelon (<i>Citrullus lanatus</i>)- Seed	South Africa	Parents plants were inspected during active growth and found to be free from a) Tobacco ringspot nepovirus b) Tomato ringspot nepovirus c) Squash mosaic comovirus
788.	Watermelon (<i>Citrullus lanatus</i>)- Seed	South Korea	Nil
789.	Watermelon (<i>Citrullus lanatus</i>)- Seed	South Korea	Nil
790.	Watermelon (<i>Citrullus lanatus</i>)- Seed	Syria	Seed is free from the following Virus relating to Watermelon: Cucumber green mottle mosaic virus, Cucumber mosaic virus, Cucumber leaf spot virus, Squash mosaic virus, Zucchini Yellow mosaic virus.
791.	Watermelon (<i>Citrullus lanatus</i>)- Seed	Tanzania	The seeds were obtained from mother plants which were inspected during active growth and found to be free from <i>Chalara elegans</i> , <i>Botryotinia fuckeliana</i> , <i>Cladosporium cucumerinum</i> , Zucchini yellow mosaic virus, <i>Rhodococcus fascians</i> and <i>pseudomonas syringae</i> pv. <i>Lachrymans</i> .
792.	Watermelon (<i>Citrullus lanatus</i>)- Seed	Uganda	Seeds have been harvested from mother plants, inspected and found to be free from following pest and diseases <i>Aphelenchoides ritizemabosi</i> , <i>alternaria brassicae</i> , <i>diaporthe phaseolorum</i> , <i>phytopathere nicotiane</i> , aster yellow phytoplasm, <i>pseudomonas syringe</i> , <i>xanthomanas vesicotoria</i> , alfalfa mosaic virus Seeds are treated with polymer and thiram
793.	Xanthosoma atrovirens (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
794.	Yard Long Bean (<i>Vigna unguiculata ssp. Sesquipedalis</i>)- Seed	Philippines	Seeds are free from soil
795.	<i>Yucca desmatiana</i> variegata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of <i>Xylella fastidiosa</i> as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....

796.	Yucca desmetiana (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
797.	Yucca gloriosa Variegata (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
798.	Zingiber officinale (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
799.	Zoysia attenuifolia (Tissue Culture)	Australia	“Prior to the removal of the plant tissue from media, the tissue cultures were inspected and found to be free of contamination. The plant tissue was aseptically transferred under supervision to sterile containers which were then sealed and not subsequently re-opened”. AND All tissue cultures in this consignment were derived from mother tissue cultures that were tested by PCR and found free of Xylella fastidiosa as indicated in laboratory test report number (TCPT14-2018). These plants are produced at their AQIS accredited lab at No.....
800.	Zuccinni (Cucurbita pepo)- seed	South Africa	Free from Tobacco ring spot nepovirus, tomato ring spot nepovirus and squash mosaic virus